arXiv:math/0005012v4 [math.AG] 6 Sep 2000

# NEF DIVISORS IN CODIMENSION ONE ON THE MODULI SPACE OF STABLE CURVES

ATSUSHI MORIWAKI

ABSTRACT. Let $M_g$ be the moduli space of smooth curves of genus $g \geq 3$, and $\bar{M}_g$ the Deligne-Mumford compactification in terms of stable curves. Let $\bar{M}_g^{[1]}$ be an open set of $\bar{M}_g$ consisting of stable curves of genus $g$ with one node at most. In this paper, we determine the necessary and sufficient condition to guarantee that a $\mathbb{Q}$-divisor $D$ on $\bar{M}_g$ is nef over $\bar{M}_g^{[1]}$, that is, $(D \cdot C) \geq 0$ for all irreducible curves $C$ on $\bar{M}_g$ with $C \cap \bar{M}_g^{[1]} \neq \emptyset$.

## CONTENTS

Introduction 1
1. Notations, conventions, terminology and preliminaries 4
2. A generalization of relative Bogomolov's inequality 12
3. A certain kind of hyperelliptic fibrations 15
4. Slope inequalities on $\bar{M}_{g,T}$ 19
5. The proof of the main result 25
References 30

## INTRODUCTION

Throughout this paper, we fix an algebraically closed field $k$, and every algebraic scheme is defined over $k$. For simplicity, we assume that the characteristic of $k$ is zero in this introduction.

Let $X$ be a normal complete variety and $\mathcal{P}$ a certain kind of positivity of $\mathbb{Q}$-line bundles on $X$ (e.g. ampleness, effectivity, bigness, etc). A problem to describe the cone $\operatorname{Cone}(X;\mathcal{P})$ consisting of $\mathbb{Q}$-line bundles with the positivity $\mathcal{P}$ is usually very hard and interesting. In this paper, as positivity, we consider numerical effectivity over a fixed open set. Namely, let $U$ be a Zariski open set of $X$. We say a $\mathbb{Q}$-line bundle $L$ is *nef over* $U$ if, for all irreducible curves $C$ with $C \cap U \neq \emptyset$, $(L \cdot C) \geq 0$. We define the relative nef cone $\operatorname{Nef}(X;U)$ over $U$ to be the cone of $\mathbb{Q}$-line bundles on $X$ which are nef over $U$.

Let $g$ and $n$ be non-negative integers with $2g - 2 + n > 0$. Let $\bar{M}_{g,n}$ (resp. $M_{g,n}$) denote the moduli space of $n$-pointed stable curves (resp. $n$-pointed smooth curves) of genus $g$. For a non-negative integer $t$, an irreducible component of the closed subscheme consisting of curves with at least $t$ nodes is called *a $t$-codimensional stratum* of $\bar{M}_{g,n}$. (For example, a $1$-codimensional stratum is a boundary component.) We denote by $S^t(\bar{M}_{g,n})$ the set of all $t$-codimensional strata of $\bar{M}_{g,n}$. Let $\bar{M}_{g,n}^{[t]}$ be the open set of $\bar{M}_{g,n}$ obtained by subtracting all $(t+1)$-codimensional strata,

---

*Date*: 6/Sep/2000, 11:50AM (JP), (Version 2.1).

i.e., $\bar{M}_{g,n}^{[t]}$ is the open set consisting of curves with at most $t$ nodes. (Note that $\bar{M}_{g,n}^{[0]} = M_{g,n}$.) Here we consider the following problem:

**Problem A.** Describe the tower of relative nef cones

$$\operatorname{Nef}(\bar{M}_{g,n}; M_{g,n}) \supseteq \operatorname{Nef}(\bar{M}_{g,n}; \bar{M}_{g,n}^{[1]}) \supseteq \cdots \supseteq \operatorname{Nef}(\bar{M}_{g,n}; \bar{M}_{g,n}^{[3g-3+n-1]}) = \operatorname{Nef}(\bar{M}_{g,n}).$$

We say a $\mathbb{Q}$-divisor on $\bar{M}_{g,n}$ is *F-nef* if the intersection number with every 1-dimensional stratum is non-negative. Let $\operatorname{FNef}(\bar{M}_{g,n})$ denote the cone consisting of F-nef $\mathbb{Q}$-divisors. Concerning the top $\operatorname{Nef}(\bar{M}_{g,n})$ of the tower, it is conjectured in [4], [7] and [5] that $\operatorname{FNef}(\bar{M}_{g,n}) = \operatorname{Nef}(\bar{M}_{g,n})$. In other words, the Mori cone of $\bar{M}_{g,n}$ is generated by 1-dimensional strata, which gives rise to a concrete description of $\operatorname{Nef}(\bar{M}_{g,n})$ (cf. [4], [7] and [5]). Moreover, it is closely related to the relative nef cone $\operatorname{Nef}(\bar{M}_{g,n}; M_{g,n})$. Actually, it was shown in [5] that if the weaker assertion $\operatorname{FNef}(\bar{M}_{g,n}) \subseteq \operatorname{Nef}(\bar{M}_{g,n}; M_{g,n})$ holds for all $g, n$, then $\operatorname{FNef}(\bar{M}_{g,n}) = \operatorname{Nef}(\bar{M}_{g,n})$. Further, as discussed in [5], $\bar{M}_{g,n}$ admits no interesting birational morphism to a projective variety. However, we can expect the rich birational geometry on $\bar{M}_{g,n}$ in terms of rational maps. In this sense, to understand the tower of relative nef cones as above might be a step toward this natural problem.

We assume that $g \geq 3$ and $n = 0$. Let $\lambda$ be the Hodge class on $\bar{M}_g$, and $\delta_{irr}, \delta_1, \ldots, \delta_{[g/2]}$ the classes of irreducible components of the boundary $\bar{M}_g \setminus M_g$. Let $\mu$ be a divisor on $\bar{M}_g$ given by

$$\mu = (8g+4)\lambda - g\delta_{irr} - \sum_{i=1}^{[g/2]} 4i(g-i)\delta_i.$$

In the paper [11], we proved that $\operatorname{Nef}(\bar{M}_g; M_g)$ is the convex hull spanned by $\mu, \delta_{irr}, \delta_1, \ldots, \delta_{[g/2]}$, that is,

$$\operatorname{Nef}(\bar{M}_g; M_g) = \mathbb{Q}_+\mu + \mathbb{Q}_+\delta_{irr} + \sum_{i=1}^{[g/2]} \mathbb{Q}_+\delta_i,$$

where $\mathbb{Q}_+ = \{x \in \mathbb{Q} \mid x \geq 0\}$. The main result of this paper is to determine $\operatorname{Nef}(\bar{M}_g; \bar{M}_g^{[1]})$.

**Theorem B** (cf. Theorem 5.1). *A $\mathbb{Q}$-divisor $a\mu + b_{irr}\delta_{irr} + \sum_{i=1}^{[g/2]} b_i\delta_i$ on $\bar{M}_g$ is nef over $\bar{M}_g^{[1]}$ if and only if the following system of inequalities hold:*

$$\begin{aligned} & a \geq \max\left\{ \frac{b_i}{4i(g-i)} \mid i = 1, \ldots, [g/2] \right\}, \\ & B_0 \geq B_1 \geq B_2 \geq \cdots \geq B_{[g/2]}, \\ & B^*_{[g/2]} \geq \cdots \geq B^*_2 \geq B^*_1 \geq B^*_0, \end{aligned}$$

*where $B_0$, $B^*_0$, $B_i$ and $B^*_i$ ($i = 1, \ldots, [g/2]$) are given by*

$$B_0 = 4b_{irr}, \quad B^*_0 = \frac{4b_{irr}}{g(2g-1)}, \quad B_i = \frac{b_i}{i(2i+1)} \quad \textit{and} \quad B^*_i = \frac{b_i}{(g-i)(2(g-i)+1)}.$$

An interesting point is that the above theorem shows us that $\mu$ is not only nef over $M_g$ but also nef over $\bar{M}_g^{[1]}$. Moreover, the theorem tells us that every nef $\mathbb{Q}$-divisor over $\bar{M}_g^{[1]}$ can be obtained in the following way. Namely, we first fix a non-negative rational number $b_{irr}$, and take $b_1$ with

$$\frac{4(g-1)b_{irr}}{g} \leq b_1 \leq 12b_{irr}.$$

Further, we choose $b_2, \dots , b_{[g/2]}$ inductively by using

$$\frac{(g-1-i)(2(g-i)-1)}{(g-i)(2(g-i)+1)} b_i \leq b_{i+1} \leq \frac{(i+1)(2i+3)}{i(2i+1)} b_i.$$

Finally, we take $a$ with

$$a \geq \max \left\{ \frac{b_i}{4i(g-i)} \mid i = 1, \dots , [g/2] \right\}.$$

Then, a $\mathbb{Q}$-divisor given by $a\mu + b_{irr}\delta_{irr} + \sum_{i=1}^{[g/2]} b_i\delta_i$ is nef over $\bar{M}_g^{[1]}$. Further, as corollaries of the above theorem, we have the following.

**Corollary C** (cf. Corollary 5.2). *For an irreducible component $\Delta$ of the boundary $\bar{M}_g \setminus M_g$, let $\widetilde{\Delta}$ be the normalization of $\Delta$, and $\rho_\Delta : \widetilde{\Delta} \to \bar{M}_g$ the induced morphism. Then, a $\mathbb{Q}$-divisor $D$ on $\bar{M}_g$ is nef over $\bar{M}_g^{[1]}$ if and only if the following are satisfied:*

(1) *$D$ is weakly positive at any points of $M_g$.*

(2) *For every boundary component $\Delta$, $\rho_\Delta^*(D)$ is weakly positive at any points of $\rho_\Delta^{-1}(\bar{M}_g^{[1]})$*

*For the definition of weak positivity, see* §1.1.

**Corollary D** (cf. Corollary 5.3). *With notation as above, if $\rho_\Delta^*(D)$ is nef over $\rho_\Delta^{-1}(\bar{M}_g^{[1]})$ for every boundary component $\Delta$, then $D$ is nef over $\bar{M}_g^{[1]}$. In particular, the Mori cone of $\bar{M}_g$ is the convex hull spanned by curves lying on the boundary $\bar{M}_g \setminus M_g$, which gives rise to a special case of* [5, Proposition 3.1].

Let us go back to the general situation. Similarly, for $\Delta \in S^l(\bar{M}_{g,n})$, let $\widetilde{\Delta}$ be the normalization of $\Delta$, and $\rho_\Delta : \widetilde{\Delta} \to \bar{M}_{g,n}$ the induced morphism. Inspired by the above corollaries, we have the following questions:

**Question E.** For a non-negative integer $t$, if a $\mathbb{Q}$-divisor $D$ on $\bar{M}_{g,n}$ is nef over $\bar{M}_{g,n}^{[t]}$, then is $\rho_\Delta^*(D)$ weakly positive at any points of $\rho_\Delta^{-1}(\bar{M}_{g,n}^{[l]})$ for all $0 \leq l \leq t$ and all $\Delta \in S^l(\bar{M}_{g,n})$? More strongly, if $D$ is nef over $\bar{M}_{g,n}^{[t]}$, then is $D$ weakly positive at any points of $\bar{M}_{g,n}^{[t]}$?

**Question F.** Fix an integer $t$ with $0 \leq t \leq 3g-3+n-1$. If $\rho_\Delta^*(D)$ is nef over $\rho_\Delta^{-1}(\bar{M}_{g,n}^{[t]})$ for all $\Delta \in S^t(\bar{M}_{g,n})$, then is $D$ is nef over $\bar{M}_{g,n}^{[t]}$?

In the case $t = 3g-3+n-1$, the above question is nothing more than asking $\mathrm{FNef}(\bar{M}_{g,n}) = \mathrm{Nef}(\bar{M}_{g,n})$.

In order to get the above theorem, we need a certain kind of slope inequalities on the moduli space of $n$-pointed stable curves. The $\mathbb{Q}$-line bundles $\lambda$ and $\psi_1, \dots , \psi_n$ on $\bar{M}_{g,n}$ are defined as follows: Let $\pi : \bar{M}_{g,n+1} \to \bar{M}_{g,n}$ be the universal curve of $\bar{M}_{g,n}$, and $s_1, \dots , s_n : \bar{M}_{g,n} \to \bar{M}_{g,n+1}$ the sections of $\pi$ arising from the $n$-points of $\bar{M}_{g,n}$. Then, $\lambda = \det(\pi_*(\omega_{\bar{M}_{g,n+1}/\bar{M}_{g,n}}))$ and $\psi_i = s_i^*(\omega_{\bar{M}_{g,n+1}/\bar{M}_{g,n}})$ for $i = 1, \dots , n$. Here we set

$$[n] = \{1, \dots , n\} \quad \text{(note that } [0] = \emptyset),$$

$$\Upsilon_{g,n} = \{(i, I) \mid i \in \mathbb{Z}, 0 \leq i \leq g \text{ and } I \subseteq [n]\} \setminus \{(0, \emptyset), (0, \{1\}), \dots , (0, \{n\})\},$$

$$\overline{\Upsilon}_{g,n} = \{\{(i, I), (j, J)\} \mid (i, I), (j, J) \in \Upsilon_{g,n}, i + j = g, I \cap J = \emptyset, I \cup J = [n]\}.$$

Moreover, for a finite set $S$, we denote the number of it by $|S|$. The boundary $\bar{M}_{g,n} \setminus M_{g,n}$ has the following irreducible decomposition:

$$\bar{M}_{g,n} \setminus M_{g,n} = \Delta_{irr} \cup \bigcup_{\{(i,I),(j,J)\} \in \overline{\Upsilon}_{g,n}} \Delta_{\{(i,I),(j,J)\}}.$$

A general point of $\Delta_{irr}$ represents an $n$-pointed irreducible stable curve with one node. A general point of $\Delta_{\{(i,I),(j,J)\}}$ represents an $n$-pointed stable curve consisting of an $|I|$-pointed smooth curve $C_1$ of genus $i$ and a $|J|$-pointed smooth curve $C_2$ of genus $j$ meeting transversally at one point, where $|I|$-points on $C_1$ (resp. $|J|$-points on $C_2$) arise from $\{s_t\}_{t \in I}$ (resp. $\{s_l\}_{l \in J}$). Let $\delta_{irr}$ and $\delta_{\{(i,I),(j,J)\}}$ be the classes of $\Delta_{irr}$ and $\Delta_{\{(i,I),(j,J)\}}$ in $\mathrm{Pic}(\bar{M}_{g,n}) \otimes \mathbb{Q}$ respectively. For a subset $L$ of $[n]$, we define a $\mathbb{Q}$-divisor $\theta_L$ on $\bar{M}_{g,n}$ to be

$$\theta_L = 4(g-1+|L|)(g-1)\sum_{t \in L} \psi_t - 12|L|^2\lambda + |L|^2\delta_{irr} - \sum_{\upsilon \in \overline{\Upsilon}_{g,n}} 4\gamma_L(\upsilon)\delta_\upsilon,$$

where $\gamma_L : \overline{\Upsilon}_{g,n} \to \mathbb{Z}$ is given by

$$\gamma_L\left(\{(i,I),(j,J)\}\right) = \left(\det\begin{pmatrix} i & |L \cap I| \\ j & |L \cap J| \end{pmatrix} + |L \cap I|\right)\left(\det\begin{pmatrix} i & |L \cap I| \\ j & |L \cap J| \end{pmatrix} - |L \cap J|\right).$$

Then, we have the following.

**Theorem G** (cf. Theorem 4.1)**.** *For any subset $L$ of $[n]$, the divisor $\theta_L$ is weakly positive at any points of $M_{g,n}$. In particular, it is nef over $M_{g,n}$.*

We remark that R. Hain has already announced the above inequality in the case where $n = 1$. (For details, see [6].) Theorem G is a generalization of his inequality.

Here we assume that $g \geq 2$. First note that

$$\mu = (8g+4)\lambda - g\delta_{irr} - \sum_{\{(i,I),(j,J)\} \in \overline{\Upsilon}_{g,n}} 4ij\delta_{\{(i,I),(j,J)\}}$$

is nef over $M_{g,n}$. Thus, as a consequence of Theorem G, we can see that

$$\mathbb{Q}_+\mu + \sum_{L \subseteq [n]} \mathbb{Q}_+\theta_L + \mathbb{Q}_+\delta_{irr} + \sum_{\upsilon \in \overline{\Upsilon}_{g,n}} \mathbb{Q}_+\delta_\upsilon \subseteq \mathrm{Nef}(\bar{M}_{g,n}; M_{g,n}),$$

so that we may ask the following question:

**Question H.** Is $\mathrm{Nef}(\bar{M}_{g,n}; M_{g,n})$ the convex hull spanned by $\mathbb{Q}$-divisors $\mu$, $\theta_L$ ($\forall L \subseteq [n]$), $\delta_{irr}$ and $\delta_\upsilon$ ($\forall \upsilon \in \overline{\Upsilon}_{g,n}$).

Corollary 4.2 and Corollary 4.3 are partial answers for the above question. If the above question is true, then it gives an affirmative answer of Question E for $t = 0$.

Finally, we would like to give hearty thanks to Prof. Hain and Prof. Keel for their useful comments for this paper.

## 1. NOTATIONS, CONVENTIONS, TERMINOLOGY AND PRELIMINARIES

Throughout this paper, we fix an algebraically closed field $k$, and every algebraic scheme is defined over $k$.

1.1. **The positivity of Weil divisors.** Let $X$ be a normal variety. Let denote $Z^1(X)$ (resp. $\operatorname{Div}(X)$) the group of Weil divisors (resp. Cartier divisors) on $X$, and $\sim$ the linear equivalence on $Z^1(X)$. We set $A^1(X) = Z^1(X)/\!\sim$ and $\operatorname{Pic}(X) = \operatorname{Div}(X)/\!\sim$. Note that $\operatorname{Pic}(X)$ is canonically isomorphic to the Picard group (the group of isomorphism classes of line bundles). Moreover, we denote by $\operatorname{Ref}(X)$ the set of isomorphism classes of reflexive sheaves of rank $1$ on $X$. For a Weil divisor $D$, the sheaf $\mathcal{O}_X(D)$ is given by

$$\mathcal{O}_X(D)(U) = \{\phi \in \operatorname{Rat}(X)^{\times} \mid (\phi) + D \text{ is effective over } U\} \cup \{0\}$$

for each Zariski open set $U$ of $X$. Then, we can see $\mathcal{O}_X(D) \in \operatorname{Ref}(X)$. Conversely, let $L$ be a reflexive sheaf of rank $1$ on $X$. For a non-zero rational section $s$ of $L$, $\operatorname{div}(s)$ is defined as follows: Let $X_0$ be the maximal Zariski open set of $X$ over which $L$ is locally free. Note that $\operatorname{codim}(X \setminus X_0) \geq 2$. Then, $\operatorname{div}(s) \in Z^1(X)$ is defined by the Zariski closure of $\operatorname{div}(\left.s\right|_{X_0})$. By our definition, we can see that $\mathcal{O}_X(\operatorname{div}(s)) \simeq L$. Thus, the correspondence $D \mapsto \mathcal{O}_X(D)$ gives rise to an isomorphism $A^1(X) \simeq \operatorname{Ref}(X)$. Here we remark that if $x \notin \operatorname{Supp}(\operatorname{div}(s))$, then $L$ is free at $x$ because $\mathcal{O}_X(\operatorname{div}(s))_x = \mathcal{O}_{X,x}$ for $x \notin \operatorname{Supp}(\operatorname{div}(s))$.

An element of $Z^1(X) \otimes \mathbb{Q}$ (resp. $\operatorname{Div}(X) \otimes \mathbb{Q}$) is celled a $\mathbb{Q}$-*divisor* (resp. $\mathbb{Q}$-*Cartier divisor*). For $\mathbb{Q}$-divisors $D_1$ and $D_2$, we say $D_1$ is $\mathbb{Q}$-linearly equivalent to $D_2$, denoted by $D_1 \sim_{\mathbb{Q}} D_2$, if there is a positive integer $n$ such that $nD_1, nD_2 \in Z^1(X)$ and $nD_1 \sim nD_2$, i.e., $D_1$ coincides with $D_2$ in $A^1(X) \otimes \mathbb{Q}$.

Fix a subset $S$ of $X$. For $D \in Z^1(X) \otimes \mathbb{Q}$, we say $D$ is *semi-ample over* $S$ if, for any $s \in S$, there is an effective $\mathbb{Q}$-divisor $E$ on $X$ with $s \notin \operatorname{Supp}(E)$ and $D \sim_{\mathbb{Q}} E$. Moreover, $D$ is said to be *weakly positive over* $S$ if there are $\mathbb{Q}$-divisors $Z_1, \dots, Z_l$, a sequence $\{D_m\}_{m=1}^{\infty}$ of $\mathbb{Q}$-divisors, and sequences $\{a_{1,m}\}_{m=1}^{\infty}, \dots, \{a_{l,m}\}_{m=1}^{\infty}$ of rational numbers such that

(1) $l$ does not depend on $m$,
(2) $D_m$ is semi-ample over $S$ for all $m \gg 0$,
(3) $D \sim_{\mathbb{Q}} D_m + \sum_{i=1}^{l} a_{i,m} Z_i$ for all $m \gg 0$, and
(4) $\lim_{m \to \infty} a_{i,m} = 0$ for all $i = 1, \dots, l$.

In the above definition, if $D$, $D_m$ and $Z_i$'s are $\mathbb{Q}$-Cartier divisors, then $D$ is said to be *weakly positive over* $S$ *in terms of Cartier divisors* (for short, C-weakly positive over $S$). Further, if $D$ is semi-ample over $\{x\}$ for some $x \in X$, then we say $D$ is semi-ample at $x$. Similarly, we define the weak positivity of $D$ at $x$ and the C-weak positivity of $D$ at $x$. We remark that weak positivity in [11] is nothing more than C-weak positivity. Moreover, note that if a $\mathbb{Q}$-divisor $D$ is semi-ample at $x$, then $D$ is a $\mathbb{Q}$-Cartier divisor around $x$, i.e., there is a Zariski open set $U$ of $X$ such that $x \in U$ and $\left.D\right|_U$ is a $\mathbb{Q}$-Cartier divisor on $U$.

A normal variety $X$ is said to be $\mathbb{Q}$-*factorial* if $Z^1(X) \otimes \mathbb{Q} = \operatorname{Div}(X) \otimes \mathbb{Q}$, i.e., any Weil divisors are $\mathbb{Q}$-Cartier divisors. It is well known that if $Y \to X$ is a finite and surjective morphism of normal varieties and $Y$ is $\mathbb{Q}$-factorial, then $X$ is also $\mathbb{Q}$-factorial (cf. [8, Lemma 5.16]). Thus the moduli space $\bar{M}_{g,n}$ of $n$-pointed stable curves of genus $g$ is $\mathbb{Q}$-factorial because $\bar{M}_{g,n}$ is an orbifold. If $X$ is $\mathbb{Q}$-factorial, then the weak positivity of $D$ over $S$ coincides with the C-weak positivity of $D$ over $S$.

We assume that $X$ is complete and $D$ is a $\mathbb{Q}$-Cartier divisor. We say $D$ is *nef over* $S$ if $(D \cdot C) \geq 0$ for any complete irreducible curves $C$ with $S \cap C \neq \emptyset$. Moreover, for a point $x$ of $X$, we say $D$ *is nef at* $x$ if $D$ is nef over $\{x\}$. Note that

$$\text{“}D \text{ is semi-ample at } x\text{”} \quad \Longrightarrow \quad \text{“}D \text{ is C-weakly positive at } x\text{”} \quad \Longrightarrow \quad \text{“}D \text{ is nef at } x\text{”}$$

**Lemma 1.1.1** ($\operatorname{char}(k) \geq 0$)**.** *Let $D$ be a $\mathbb{Q}$-divisor on $X$, and $x_1, \ldots, x_n \in X$. If $D$ is semi-ample at $x_i$ for each $i$, then there is an effective $\mathbb{Q}$-divisor $E$ on $X$ such that $E \sim_{\mathbb{Q}} D$ and $x_i \notin \operatorname{Supp}(E)$ for all $i$.*

*Proof.* By our assumption, there is an effective $\mathbb{Q}$-divisor $E_i$ on $X$ such that $E_i \sim_{\mathbb{Q}} D$ and $x_i \notin \operatorname{Supp}(E_i)$. Take a sufficiently large integer $m$ such that $mD, mE_1, \ldots, mE_n \in Z^1(X)$ and $mD \sim mE_i$ for all $i$. Thus, there is a section $s_i$ of $H^0(X, \mathcal{O}_X(mD))$ with $\operatorname{div}(s_i) = mE_i$. Here since $x_i \notin \operatorname{Supp}(mE_i)$ and $\mathcal{O}_X(mD) \simeq \mathcal{O}_X(mE_i)$, we can see that $\mathcal{O}_X(mD)$ is free at each $x_i$.

For $\alpha = (\alpha_1, \ldots \alpha_n) \in k^n$, we set $s_\alpha = \alpha_1 s_1 + \cdots + \alpha_n s_n \in H^0(X, \mathcal{O}_X(mD))$. Further, we set $V_i = \{\alpha \in k^n \mid s_\alpha(x_i) = 0\}$. Then, $\dim V_i = n - 1$ for all $i$. Thus, since $\#(k) = \infty$, there is $\alpha \in k^n$ with $\alpha \notin V_1 \cup \cdots \cup V_r$, i.e., $s_\alpha(x_i) \neq 0$ for all $i$. Let us consider a divisor $E = \operatorname{div}(s_\alpha)$. Then, $E \sim mD$ and $x_i \notin \operatorname{Supp}(E)$ for all $i$. □

**Proposition 1.1.2** ($\operatorname{char}(k) \geq 0$)**.** *Let $\pi : X \to Y$ be a surjective, proper and generically finite morphism of normal varieties. Let $D$ be a $\mathbb{Q}$-divisor on $X$ and $S$ a subset of $Y$ such that $\pi^{-1}(S)$ is finite. Then, we have the following.*

(1) *If $D$ is semi-ample over $\pi^{-1}(S)$, then $\pi_*(D)$ is semi-ample over $S$.*
(2) *If $D$ is weakly positive over $\pi^{-1}(S)$, then $\pi_*(D)$ is weakly positive over $S$.*

*Proof.* (1) By Lemma 1.1.1, there is an effective divisor $E$ on $X$ such that $E \sim_{\mathbb{Q}} D$ and $s' \notin \operatorname{Supp}(E)$ for all $s' \in \pi^{-1}(S)$. Then, $\pi_*(E) \sim_{\mathbb{Q}} \pi_*(D)$ and $s \notin \pi(\operatorname{Supp}(E)) = \operatorname{Supp}(\pi_*(E))$ for all $s \in S$.

(2) This is a consequence of (1). □

**Proposition 1.1.3** ($\operatorname{char}(k) \geq 0$)**.** *Let $\pi : X \to Y$ be a surjective, proper morphism of normal varieties. We assume that $Y$ is $\mathbb{Q}$-factorial. Let $D$ be a $\mathbb{Q}$-divisor on $Y$, and $S$ a subset of $Y$. Then, we have the following.*

(1) *If $D$ is semi-ample over $S$, then $\pi^*(D)$ is semi-ample over $f^{-1}(S)$.*
(2) *If $D$ is weakly positive over $S$, then $\pi^*(D)$ is C-weakly positive over $S$.*

*Proof.* (1) Let $s'$ be a point in $\pi^{-1}(S)$. Then, there is an effective $\mathbb{Q}$-divisor $E$ on $Y$ with $D \sim_{\mathbb{Q}} E$ and $\pi(s') \notin \operatorname{Supp}(E)$. Thus, $\pi^*(D) \sim_{\mathbb{Q}} \pi^*(E)$ and $s' \notin \operatorname{Supp}(\pi^*(E))$. Therefore, $\pi^*(D)$ is semiample over $\pi^{-1}(S)$.

(2) This is a consequence of (1). □

**Lemma 1.1.4** ($\operatorname{char}(k) \geq 0$)**.** *Let $X$ and $Y$ be complete varieties, and let $D$ and $E$ be $\mathbb{Q}$-Cartier divisors on $X$ and $Y$ respectively. Let $p : X \times Y$ and $q : X \times Y \to Y$ be the projections to the first factor and the second factor respectively. For $(x, y) \in X \times Y$, $p^*(D) + q^*(E)$ is nef at $(x, y)$ if and only if $D$ and $E$ are nef at $x$ and $y$ respectively.*

*Proof.* First we assume that $p^*(D) + q^*(E)$ is nef at $(x, y)$. Let $C$ be a complete irreducible curve on $X$ with $x \in C$. Then, $C_y = C \times \{y\}$ is a complete curve on $X \times Y$ with $(x, y) \in C_y$. Moreover, $(p^*(D) + q^*(E) \cdot C_y) = (D \cdot C)$. Thus, $(D \cdot C) \geq 0$, which says us that $D$ is nef at $x$. In the same way, we can see that $E$ is nef at $y$.

Next we assume that $D$ and $E$ are nef at $x$ and $y$ respectively. In order to see that $p^*(D) + q^*(E)$ is nef at $(x, y)$, it is sufficient to check that $(p^*(D) \cdot C) \geq 0$ and $(q^*(E) \cdot C) \geq 0$ for any complete irreducible curves $C$ on $X \times Y$ with $(x, y) \in C$. Here, $p(C)$ is either $\{x\}$, or a complete irreducible curve passing through $x$. Thus, by virtue of the projection formula, $(p^*(D)\cdot C)\geq 0$. In the same way, $(q^*(E)\cdot C)\geq 0$. $\square$

1.2. **The first Chern class of coherent sheaves.** Let $X$ be a normal variety, and $F$ a coherent $\mathcal{O}_X$-module on $X$. Here we define $c_1(F)\in A^1(X)$ in the following way.

Case 1. $F$ is a torsion sheaf. In this case, we set

$$D=\sum_{\substack{P\in X,\\ \operatorname{depth}(P)=1}}\operatorname{lenght}(F_P)\overline{\{P\}},$$

where $\overline{\{P\}}$ is the Zariski closure of $\{P\}$ in $X$. Then, $c_1(F)$ is defined by the class of $D$.

Case 2. $F$ is a torsion free sheaf. Let $r$ be the rank of $F$. Then, $(\bigwedge^r F)^{\vee\vee}$ is a reflexive sheaf of rank 1, where ${}^{\vee\vee}$ means the double dual of sheaves. Thus, we define $c_1(F)$ to be the class of $(\bigwedge^r F)^{\vee\vee}$.

Case 3. $F$ is general. Let $T$ be the torsion part of $F$. Then, $c_1(F)=c_1(T)+c_1(F/T)$.

Note that if $0\to F_1\to F_2\to F_3\to 0$ is an exact sequence of coherent $\mathcal{O}_X$-modules, then $c_1(F_2)=c_1(F_1)+c_1(F_3)$. Moreover, let $L$ be a reflexive sheaf of rank 1 on $X$, and $s$ a non-zero section of $L$. Then

$$c_1(L)=c_1\left(\operatorname{Coker}(\mathcal{O}_X\xrightarrow{\times s}L)\right)=\text{the class of }\operatorname{div}(s).$$

**Proposition 1.2.1** ($\operatorname{char}(k)\geq 0$)**.** *Let $X$ be a normal algebraic variety, $F$ a coherent $\mathcal{O}_X$-module, and $x$ a point of $X$. If $F$ is generated by global sections at $x$ and $F$ is free at $x$, then $c_1(F)$ is semi-ample at $x$.*

*Proof.* Let $T$ be the torsion part of $F$. Then, $c_1(F)=c_1(F/T)+c_1(T)$. Here since $F$ is free at $x$, $c_1(T)$ is semi-ample at $x$. Moreover, it is easy to see that $F/T$ is generated by global sections at $x$. Therefore, to prove our proposition, we may assume that $F$ is a torsion free sheaf.

Let $r$ be the rank of $F$ and $\kappa(x)$ the residue field of $x$. Then, by our assumption, there are sections $s_1,\ldots,s_r$ of $F$ such that $\{s_i(x)\}$ forms a basis of $F\otimes\kappa(x)$. Since we can view $s_i$ as an injection $s_i:\mathcal{O}_X\to F$, $s=s_1\wedge\cdots\wedge s_r$ gives rise to an injection $s:\mathcal{O}_X\to(\bigwedge^r F)^{\vee\vee}$, which is bijective at $x$. Thus, $x\notin\operatorname{div}(s)$. $\square$

1.3. **The discriminant divisor of vector bundles.** Let $f:X\to Y$ be a proper surjective morphism of algebraic varieties of the relative dimension one, and let $E$ be a locally free sheaf on $X$. We define the *discriminant divisor of $E$ with respect to $f$* to be

$$\operatorname{dis}_{X/Y}(E)=f_*\left(2\operatorname{rk}(E)c_2(E)-(\operatorname{rk}(E)-1)c_1(E)^2\right).$$

**Lemma 1.3.1** ($\operatorname{char}(k)\geq 0$)**.** *Let $f:X\to Y$ be a flat, surjective and projective morphism of varieties with $\dim f=1$. Let $E$ be a vector bundle of rank $r$ on $X$. Then, we have the following.*

(1) $\operatorname{dis}_{X/Y}(E)$ *is a Cartier divisor.*

(2) *Let* $u : Y' \to Y$ *be a morphism of varieties, and let*

$$\begin{tikzcd} X \arrow[d, "f"'] & X \times_Y Y' \arrow[l, "u'"'] \arrow[d, "f'"] \\ Y & Y' \arrow[l, "u"'] \end{tikzcd}$$

*be the induced diagram of the fiber product. If* $X \times_Y Y'$ *is integral, then* $\operatorname{dis}_{X\times_Y Y'/Y'}(u'^*(E)) = u^*(\operatorname{dis}_{X/Y}(E))$.

*Proof.* (1) We set $F = \mathcal{E}nd(E)$. Let $p : P = \mathbb{P}(F) \to X$ be the projective bundle of $F$, and $\mathcal{O}_P(1)$ the tautological line bundle on $P$. Let $g : P \to Y$ be the composition of $P \xrightarrow{p} X \xrightarrow{f} Y$. Then, since

$$p_*(c_1(\mathcal{O}_P(1))^{r^2+1}) = -c_2(F) = -(2rc_2(E) - (r-1)c_1(E)^2),$$

we have $g_*(c_1(\mathcal{O}_P(1))^{r^2+1}) = -\operatorname{dis}_{X/Y}(E)$. Thus, $\operatorname{dis}_{X/Y}(E) = -c_1\left(\langle \mathcal{O}_P(1)^{\cdot r^2+1}\rangle(P/Y)\right)$, where

$$\langle\ ,\cdots,\ \rangle(P/Y) : \overbrace{\operatorname{Pic}(P) \times \cdots \times \operatorname{Pic}(P)}^{\dim g+1} \to \operatorname{Pic}(Y)$$

is Deligne's pairing for the flat morphism $g : P \to Y$. Therefore, $\operatorname{dis}_{X/Y}(E)$ is a Cartier divisor.

(2) This follows from the compatibility of Deligne's pairing by base changes. □

**Remark 1.3.2.** In (2) of Lemma 1.3.1, $X \times_Y Y'$ is integral if the generic fiber of $X \times_Y Y' \to Y'$ is integral by virtue of [12, Lemma 4.2].

1.4. **The moduli space of $T$-pointed stable curves of genus $g$.** Let $g$ be a non-negative integer and $T$ a finite set with $2g-2+|T| > 0$, where $|T|$ is the number of $T$. Recall that $[n] = \{1, \ldots, n\}$ and $[0] = \emptyset$. Usually, we use $[n]$ as $T$. Let $\bar{M}_{g,T}$ (resp. $M_{g,T}$) denote the moduli space of $T$-pointed stable curves (resp. $T$-pointed smooth curves) of genus $g$, namely, $\bar{M}_{g,T}$ (resp. $M_{g,T}$) is the moduli space of $|T|$-pointed stable curves (resp. $|T|$-pointed smooth curves) of genus $g$, whose marked points are labeled by the index set $T$.

Roughly speaking, the $\mathbb{Q}$-line bundles $\lambda$ and $\{\psi_t\}_{t\in T}$ on $\bar{M}_{g,T}$ are defined as follows: Let $\pi : \mathcal{C} \to \bar{M}_{g,T}$ be the universal curve of $\bar{M}_{g,T}$, and $s_t : \bar{M}_{g,T} \to \mathcal{C}$ ($t \in T$) the sections of $\pi$ arising from the $T$-points of $\bar{M}_{g,T}$. Then, $\lambda = \det(\pi_*(\omega_{\mathcal{C}/\bar{M}_{g,T}}))$ and $\psi_t = s_t^*(\omega_{\mathcal{C}/\bar{M}_{g,T}})$ for $t \in T$.

For $x \in \bar{M}_{g,T}$, let denote $C_x$ the nodal curve corresponding to $x$ (here we forget the $T$-points). Let $S^l(\bar{M}_{g,T})$ be the set of all irreducible components of the closed set

$$\{x \in \bar{M}_{g,T} \mid \#(\operatorname{Sing}(C_x)) \geq l\}.$$

Then, every element of $S^l(\bar{M}_{g,T})$ is of codimension $l$, so that it is called an *$l$-codimensional stratum* of $\bar{M}_{g,T}$. Note that $\bar{M}_{g,T} \setminus M_{g,T}$ is a normal crossing divisor in the sense of orbifolds. Thus the normalization of an element of $S^l(\bar{M}_{g,T})$ is $\mathbb{Q}$-factorial. Moreover, we set

$$\bar{M}_{g,T}^{[l]} = \bar{M}_{g,T} \setminus \bigcup_{\Delta \in S^{l+1}(\bar{M}_{g,T})} \Delta,$$

i.e., $\bar{M}_{g,T}^{[l]} = \{x \in \bar{M}_{g,T} \mid \#(\operatorname{Sing}(C_x)) \leq l\}$. Note that $\bar{M}_{g,T}^{[0]} = M_{g,T}$.

To describe the boundary of $\bar{M}_{g,T}$, we set

$$\begin{aligned}\Upsilon_{g,T} &= \{(i,I) \mid i \in \mathbb{Z}, 0 \le i \le g \text{ and } I \subseteq T\} \setminus (\{(0,\emptyset)\} \cup \{(0,\{t\})\}_{t\in T}),\\ \overline{\Upsilon}_{g,T} &= \{\{(i,I),(j,J)\} \mid (i,I),(j,J) \in \Upsilon_{g,T}, i+j=g, I\cap J=\emptyset, I\cup J = T\}.\end{aligned}$$

Then, the boundary $\Delta = \bar{M}_{g,T} \setminus M_{g,T}$ has the following irreducible decomposition:

$$\Delta = \Delta_{irr} \cup \bigcup_{\{(i,I),(j,J)\}\in\overline{\Upsilon}_{g,T}} \Delta_{\{(i,I),(j,J)\}}.$$

A general point of $\Delta_{irr}$ represents a $T$-pointed irreducible stable curve with one node. A general point of $\Delta_{\{(i,I),(j,J)\}}$ represents a $T$-pointed stable curve consisting of an $I$-pointed smooth curve of genus $i$ and a $J$-pointed smooth curve of genus $j$ meeting transversally at one point.



$\Delta_{irr}$



$\Delta_{\{(i,I),(j,J)\}}$

Let $\delta_{irr}$ and $\delta_{\{(i,I),(j,J)\}}$ be the classes of $\Delta_{irr}$ and $\Delta_{\{(i,I),(j,J)\}}$ in $\operatorname{Pic}(\bar{M}_{g,T}) \otimes \mathbb{Q}$ respectively. Strictly speaking, $\delta_{irr} = c_1(\mathcal{O}_{\bar{M}_{g,T}}(\Delta_{irr}))$ and

$$\delta_\upsilon = \begin{cases} \dfrac{1}{2} c_1\left(\mathcal{O}_{\bar{M}_{g,T}}(\Delta_\upsilon)\right) & \text{if } \upsilon = \{(1,\emptyset),(g-1,T)\},\\ \\ c_1\left(\mathcal{O}_{\bar{M}_{g,T}}(\Delta_\upsilon)\right) & \text{otherwise}.\end{cases}$$

In the case where $T=\emptyset$, we denote $\delta_{\{(i,\emptyset),(j,\emptyset)\}}$ by $\delta_{\{i,j\}}$ or $\delta_{\min\{i,j\}}$, i.e.,

$$\delta_i = \delta_{\{i,g-i\}} = \delta_{\{(i,\emptyset),(g-i,\emptyset)\}} \qquad (i = 1, \ldots, [g/2])$$

on $\bar{M}_g$.

Let $(Z; \{P_t\}_{t\in T})$ be a $T$-pointed stable curve of genus $g$ over $k$. Let $Q$ be a node of $Z$, and $Z_Q$ the partial normalization of $Z$ at $Q$. Then, the type of $Q$ is defined as follows:

- The case where $Z_Q$ is connected. Then, $Q$ is of type $0$.
- The case where $Z_Q$ is not connected. Let $Z_1$ and $Z_2$ be two connected components of $Z_Q$. Let $i$ (resp. $j$) be the arithmetic genus of $Z_1$ (resp. $Z_2$). Let $I = \{t \in T \mid P_t \in Z_1\}$ and $J = \{t \in T \mid P_t \in Z_2\}$. Then, we say $Q$ is of type $\{(i,I),(j,J)\}$.

In the case where $T=\emptyset$, for simplicity, a node of type $\{(i,\emptyset),(j,\emptyset)\}$ is said to be of type $i$, where $i \le j$.

Let $Y$ be a normal variety, and let $f : X \to Y$ be a $T$-pointed stable curve of genus $g$ over $Y$. Let $Y_0$ be the maximal open set over which $f$ is smooth. Assume that $Y_0 \neq \emptyset$. For $x \in X$, we define $\operatorname{mult}_x(X)$ to be $\operatorname{length}_{\mathcal{O}_{X,x}}(\omega_{X/Y}/\Omega_{X/Y})$. If $x$ is the generic point of a subvariety $T$, then we denote $\operatorname{mult}_x(X)$ by $\operatorname{mult}_T(X)$. If $x$ is closed, $Y$ is smooth at $f(x)$ and $Y \setminus Y_0$ is smooth at $f(x)$, then $X$ is locally given by $\{xy = t^{\operatorname{mult}_x(X)}\}$ around $x$, where $t$ is a defining equation of $Y \setminus Y_0$ around $f(x)$. Thus, if $Y$ is a curve, then the type of singularity at $x$ is $A_{\operatorname{mult}_x(X)-1}$.

Here, for $\upsilon \in \overline{\Upsilon}_{g,T}$, let $S(X/Y)_{\upsilon}$ (resp. $S(X/Y)_{irr}$) be the set of irreducible components of $\operatorname{Sing}(f)$ such that the type of $s$ in $f^{-1}(f(s))$ for a general $s \in S(X/Y)_{\upsilon}$ (resp. $S(X/Y)_{irr}$) is $\upsilon$ (resp. $0$). We set

$$\delta_{\upsilon}(X/Y) = \sum_{S \in S(X/Y)_{\upsilon}} \operatorname{mult}_S(X) f_*(S)$$

and

$$\delta_{irr}(X/Y) = \sum_{S \in S(X/Y)_{irr}} \operatorname{mult}_S(X) f_*(S).$$

Then, $\delta_{irr}$ and $\delta_{\upsilon}$ are normalized to guarantee the following formula:

$$\delta_{irr}(X/Y) = \varphi^*(\delta_{irr}) \quad \text{and} \quad \delta_{\upsilon}(X/Y) = \varphi^*(\delta_{\upsilon})$$

in $A^1(Y) \otimes \mathbb{Q}$, where $\varphi : Y \to \bar{M}_{g,T}$ is the induced morphism by $X \to Y$.

1.5. **The clutching maps.** Here let us consider the clutching maps and their properties.

Let $\pi : X \to Y$ be a prestable curve, i.e., $\pi : X \to Y$ is a flat and proper morphism such that the geometric fibers of $\pi$ are reduced curves with at most ordinary double points. We don't assume the connectedness of fibers. Let $s_1, s_2 : Y \to X$ be two non-crossing sections such that $\pi$ is smooth at points $s_1(y)$ and $s_2(y)$ ($\forall y \in Y$). Then, by virtue of [9, Theorem 3.4], we have the clutching diagram:

$$\begin{tikzcd}
X \arrow[rr, "p"] \arrow[dr, "\pi"] & & X' \arrow[dl, "\pi'"'] \\
& Y \arrow[ul, bend left=20, "s_1"'] \arrow[ul, bend left=50, "s_2"] \arrow[ur, bend right=30, "s"'] &
\end{tikzcd}$$

Roughly speaking, $X'$ is a prestable curve over $Y$ obtained by identifying $s_1(Y)$ with $s_2(Y)$, and $s$ is a section of $X' \to Y$ with $p \cdot s_1 = p \cdot s_2 = s$. For details, see [9, Theorem 3.4].

We assume that $\pi' : X' \to Y$ is a $T$-pointed stable curve of genus $g$, and $s$ is one of sections of $\pi' : X' \to Y$ arising from $T$-points of $\pi' : X' \to Y$. Let $\varphi : Y \to \bar{M}_{g,T}$ be the induced morphism. Here we set $\Lambda = \det(R\pi_*(\omega_{X/Y}))$, $\Delta = \det(R\pi_*(\omega_{X/Y}/\Omega_{X/Y}))$ and $\Psi = s_1^*(\omega_{X/Y}) \otimes s_2^*(\omega_{X/Y})$. Then, we have the following.

**Proposition 1.5.1.** *For simplicity, the divisor $\delta_{irr}$ on $\bar{M}_{g,T}$ is denoted by $\delta_0$.*

1. $\varphi^*(\lambda) = \Lambda$ *and* $\varphi^*(\delta) = -\Psi + \Delta$, *where* $\delta = \delta_0 + \sum_{\upsilon \in \overline{\Upsilon}_{g,T}} \delta_{\upsilon}$.
2. *We assume that* $\pi(\operatorname{Sing}(\pi)) \neq Y$ *and every geometric fiber of* $\pi$ *has one node at most. Let*

$$\Delta = \Delta_0 + \sum_{\upsilon \in \overline{\Upsilon}_{g,T}} \Delta_{\upsilon}$$

*be the decomposition such that the node of* $\pi^{-1}(x)$ ($x \in (\Delta_t)_{red}$) *gives rise to a node of type* $t$ *in* ${\pi'}^{-1}(x)$. *Moreover, let* $a$ *be the type of* $s(y)$ *in* ${\pi'}^{-1}(y)$ ($y \in Y$). *Then,*

$$\varphi^*(\delta_t) = \begin{cases} -\Psi + \Delta_a & \textit{if } t = a \\ \Delta_t & \textit{if } t \neq a \end{cases}$$

*Proof.* (1) Since $\det(R{\pi'}_*(\omega_{X'/Y})) = \det(R\pi_*(\omega_{X/Y}))$, the first statement is obvious. Thus, we can see that

$$\begin{aligned}\varphi^*(\delta) &= \det(R{\pi'}_*(\omega_{X'/Y}/\Omega_{X'/Y})) \\ &= \det(R{\pi'}_*(\omega_{X'/Y})) - \det(R{\pi'}_*(\Omega_{X'/Y})) \\ &= \Lambda - \det(R{\pi'}_*(\Omega_{X'/Y})).\end{aligned}$$

On the other hand, by [9, Theorem 3.5], there is an exact sequence

$$0 \to s^*(\Psi) \to \Omega_{X'/Y} \to p_*(\Omega_{X/Y}) \to 0.$$

Therefore, we get (1).

(2) This is a consequence of (1). □

As a corollary, we have the following.

**Corollary 1.5.2.** (1) *Let $a$ and $b$ be non-negative integers, and $T$ and $S$ non-empty finite sets with $T \cap S = \emptyset$, $2a - 2 + |T| > 0$ and $2b - 2 + |S| > 0$. Let us fix $s \in S$ and $t \in T$, and set $T' = T \setminus \{t\}$ and $S' = S \setminus \{s\}$. Let $\alpha : \bar{M}_{a,T} \times \bar{M}_{b,B} \to \bar{M}_{a+b,T' \cup S'}$ be the clutching map, and $p : \bar{M}_{a,T} \times \bar{M}_{b,S} \to \bar{M}_{a,T}$ and $q : \bar{M}_{a,T} \times \bar{M}_{b,S} \to \bar{M}_{b,S}$ the projection to the first factor and the projection to the second factor respectively. We set divisors $D \in \mathrm{Pic}(\bar{M}_{a+b,T'\cup S'}) \otimes \mathbb{Q}$, $E \in \mathrm{Pic}(\bar{M}_{a,T}) \otimes \mathbb{Q}$ and $F \in \mathrm{Pic}(\bar{M}_{b,S}) \otimes \mathbb{Q}$ as follows:*

$$\begin{aligned}D &= c\lambda + \sum_{l \in T' \cup S'} d_l \psi_l + e_{irr}\delta_{irr} + \sum_{\{(i,I),(j,J)\} \in \overline{\Upsilon}_{a+b,T'\cup S'}} e_{\{(i,I),(j,J)\}}\delta_{\{(i,I),(j,J)\}}, \\ E &= c\lambda - e_{\{(a,T'),(b,S')\}}\psi_t + \sum_{l \in T'} d_l\psi_l + e_{irr}\delta_{irr} \\ &\qquad + \sum_{\substack{\{(i',I'),(j',J')\} \in \overline{\Upsilon}_{a,T} \\ t \in J'}} e_{\{(i',I'),(j'+b,J'\cup S'\setminus\{t\})\}}\delta_{\{(i',I'),(j',J')\}}, \\ F &= c\lambda - e_{\{(a,T'),(b,S')\}}\psi_s + \sum_{l \in S'} d_l\psi_l + e_{irr}\delta_{irr} \\ &\qquad + \sum_{\substack{\{(i'',I''),(j'',J'')\} \in \overline{\Upsilon}_{b,S} \\ s \in J''}} e_{\{(i'',I''),(j''+a,J''\cup T'\setminus\{s\})\}}\delta_{\{(i'',I''),(j'',J'')\}}.\end{aligned}$$

*Then $\alpha^*(D) = p^*(E) + q^*(F)$.*

(2) *Let $g$ be a non-negative integer and $T$ a finite set with $|T| \geq 2$ and $2g - 2 + |T| > 0$. Let us fix two elements $t, t' \in T$, and set $T' = T \setminus \{t, t'\}$. Let $\beta : \bar{M}_{g,T} \to \bar{M}_{g+1,T'}$ be the clutching*

*map. We set* $D \in \operatorname{Pic}(\bar{M}_{g+1,T'}) \otimes \mathbb{Q}$ *and* $E \in \operatorname{Pic}(\bar{M}_{g,T}) \otimes \mathbb{Q}$ *as follows:*

$$D = c\lambda + \sum_{l \in T'} d_l \psi_l + e_{irr}\delta_{irr} + \sum_{\{(i,I),(j,J)\} \in \overline{\Upsilon}_{g+1,T'}} e_{\{(i,I),(j,J)\}}\delta_{\{(i,I),(j,J)\}},$$

$$\begin{aligned} E = c\lambda - e_{irr}(\psi_t + \psi_{t'}) + \sum_{l \in T'} d_l \psi_l + e_{irr}\delta_{irr} + \sum_{\substack{\{(i',I'),(j',J')\} \in \overline{\Upsilon}_{g,T} \\ t \in I', t' \in J'}} e_{irr}\delta_{\{(i',I'),(j',J')\}} \\ + \sum_{\substack{\{(i',I'),(j',J')\} \in \overline{\Upsilon}_{g,T} \\ t, t' \in J'}} e_{\{(i',I'),(j'+1,J' \setminus \{t,t'\})\}}\delta_{\{(i',I'),(j',J')\}}. \end{aligned}$$

*Then* $\beta^*(D) = E$.

*Proof.* In the following, for $x \in \bar{M}_{*,*}$, we denote by $C_x$ the corresponding nodal curve to $x$.

(1) If $C_{\alpha(x,y)}$ has two nodes, then we denote by $ty(x,y)$ the type of the node different from the node arising from the clutching map. Then,

$$ty(x,y) = \begin{cases} \{(i',I'),(j'+b, J' \cup S' \setminus \{t\})\} & \text{if } x \in \Delta_{\{(i',I'),(j',J')\}} \cap \bar{M}^{[1]}_{a,T},\ y \in M_{b,S} \text{ and } t \in J', \\ \{(i'',I''),(j''+a, J'' \cup T' \setminus \{s\})\} & \text{if } x \in M_{a,T},\ y \in \Delta_{\{(i'',I''),(j'',J'')\}} \cap \bar{M}^{[1]}_{b,S} \text{ and } s \in J''. \end{cases}$$

Thus, we get (1) by the above proposition.

(2) In the same way as above, if $C_{\beta(x)}$ has two nodes, then we denote by $ty'(x)$ the type of the node different from the node arising from the clutching map. Then,

$$ty'(x) = \begin{cases} 0 & \text{if } x \in \left(\Delta_{irr} \cup \bigcup_{t \in I', t' \in J'} \Delta_{\{(i',I'),(j',J')\}}\right) \cap \bar{M}^{[1]}_{g,T}, \\ \{(i',I'),(j'+1, J' \setminus \{t,t'\})\} & \text{if } x \in \Delta_{\{(i',I'),(j',J')\}} \cap \bar{M}^{[1]}_{g,T} \text{ and } t, t' \in J', \end{cases}$$

which implies (2) by the above proposition. □

## 2. A GENERALIZATION OF RELATIVE BOGOMOLOV'S INEQUALITY

Let $f : X \to Y$ be a projective morphism of quasi-projective varieties of the relative dimension one, and let $E$ be a locally free sheaf on $X$. Let us fix a point $y \in Y$. Assume that $f$ is smooth over $y$ and $E|_{f^{-1}(y)}$ is strongly semistable. In the paper [11], we proved that $\operatorname{dis}_{X/Y}(E)$ is weakly positive at $y$ under the assumption that $Y$ is smooth. In this section, we generalize it to the case where $Y$ is normal.

**Proposition 2.1** ($\operatorname{char}(k) \geq 0$)**.** *Let* $X$ *and* $Y$ *be algebraic varieties, and* $f : X \to Y$ *a surjective and projective morphism of* $\dim f = d$. *Let* $L$ *and* $A$ *be line bundles on* $X$. *If* $Y$ *is normal, then there are* $\mathbb{Q}$*-divisors* $Z_0, \ldots, Z_d$ *on* $Y$ *such that*

$$c_1\left(Rf_*(L^{\otimes n} \otimes A)\right) \sim_{\mathbb{Q}} \frac{f_*(c_1(L)^{d+1})}{(d+1)!} n^{d+1} + \sum_{i=0}^{d} Z_i n^i$$

*for all* $n > 0$.

*Proof.* We set $Y^0 = Y \setminus \operatorname{Sing}(Y)$, $X^0 = f^{-1}(Y^0)$ and $f^0 = f|_{X^0}$. Then, we have

$$c_1\left(Rf^0_*\left((L^{\otimes n} \otimes A)\big|_{X^0}\right)\right) = c_1(Rf_*(L^{\otimes n} \otimes A))\big|_{Y^0}$$

and

$$f^0_*\left(c_1\left(L|_{X^0}\right)^{d+1}\right) = f_*(c_1(L)^{d+1})\big|_{Y^0}.$$

Thus, by virtue of [11, Lemma 2.3], there are $\mathbb{Q}$-divisors $Z^0_0, \ldots, Z^0_d$ on $Y^0$ such that

$$c_1(Rf_*(L^{\otimes n} \otimes A))\big|_{Y^0} \sim_{\mathbb{Q}} \frac{f_*(c_1(L)^{d+1})\big|_{Y^0}}{(d+1)!} n^{d+1} + \sum_{i=0}^{d} Z^0_i n^i$$

for all $n > 0$. Let $Z_i$ be the Zariski closure of $Z^0_i$ in $Y$. Then, since $\operatorname{codim}(\operatorname{Sing}(Y)) \geq 2$,

$$c_1(Rf_*(L^{\otimes n} \otimes A)) \sim_{\mathbb{Q}} \frac{f_*(c_1(L)^{d+1})}{(d+1)!} n^{d+1} + \sum_{i=0}^{d} Z_i n^i$$

for all $n > 0$. □

**Theorem 2.2** ($\operatorname{char}(k) \geq 0$)**.** *Let $X$ be a quasi-projective variety, $Y$ a normal quasi-projective variety, and $f : X \to Y$ a surjective and projective morphism of $\dim f = 1$. Let $F$ be a locally free sheaf on $X$ with $f_*(c_1(F)) = 0$, and $S$ a finite subset of $Y$. We assume that $f$ is flat over any points of $S$, and that, for all $s \in S$, there are line bundles $L_{\bar{s}}$ and $M_{\bar{s}}$ on the geometric fiber $X_{\bar{s}}$ over $s$ such that*

$$H^0(X_{\bar{s}}, \operatorname{Sym}^m(F_{\bar{s}}) \otimes L_{\bar{s}}) = H^1(X_{\bar{s}}, \operatorname{Sym}^m(F_{\bar{s}}) \otimes M_{\bar{s}}) = 0$$

*for $m \gg 0$. Then, $f_*\left(c_2(F) - c_1(F)^2\right)$ is weakly positive over $S$.*

*Proof.* The proof of this theorem is exactly same as [11, Theorem 2.4] using Proposition 2.1, Proposition 1.2.1 and [11, Proposition 2.2]. For reader's convenience, we give the sketch of the proof of it.

Let $A$ be a very ample line bundle on $X$ such that $A_{\bar{s}} \otimes L_{\bar{s}}$ and $A_{\bar{s}} \otimes M_{\bar{s}}^{\otimes -1}$ are very ample on $X_{\bar{s}}$ for all $s \in S$. Then, we can see the following claim in the same way as in [11, Claim 2.4.1]

**Claim 2.2.1.** *$H^0(X_s, \operatorname{Sym}^m(F_s) \otimes A_s^{\otimes -1}) = H^1(X_s, \operatorname{Sym}^m(F_s) \otimes A_s) = 0$ for all $s \in S$ and $m \gg 0$.*

Since $X$ is an integral scheme of dimension greater than or equal to 2, and $X_s$ ($s \in S$) is a 1-dimensional scheme over $\kappa(s)$, there is $B \in |A^{\otimes 2}|$ such that $B$ is integral, and that $B \cap X_s$ is finite for all $s \in S$, i.e., $B$ is finite over any points of $S$. Let $\pi : B \to Y$ be the morphism induced by $f$. Let $H$ be an ample line bundle on $Y$ such that $\pi_*(F_B) \otimes H$ and $\pi_*(A_B) \otimes H$ are generated by global sections at any points of $S$, where $F_B = F|_B$ and $A_B = A|_B$.

By using Proposition 2.1, there are $\mathbb{Q}$-divisors $Z_0, \ldots, Z_r$ on $Y$ such that

$$\sum_{i \geq 0} (-1)^i c_1\left(R^i f_*\left(\operatorname{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right)\right)$$

$$\sim_{\mathbb{Q}} -\frac{1}{(r+1)!} f_*(c_2(F) - c_1(F)^2) m^{r+1} + \sum_{i=0}^{r} Z_i m^i$$

in the same way as in the proof of [11, Theorem 2.4]. The following claim also can be proved in the same way as in [11, Claim 2.4.2].

**Claim 2.2.2.** *If $m \gg 0$, then we have the following.*

(a) $c_1\left(R^i f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right)\right) = 0$ *for all* $i \geq 2$.
(b) $f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right) = 0$.
(c) $R^1 f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right)$ *is free at any points of* $S$.
(d) $R^1 f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A \otimes f^*(H)\right) = 0$ *around any points of* $S$.

By (a) and (b) of Claim 2.2.2,

$$\frac{f_*(c_2(F) - c_1(F)^2)}{(r+1)!} \sim_{\mathbb{Q}} \frac{c_1\left(R^1 f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right)\right)}{m^{r+1}} + \sum_{i=0}^{r} \frac{Z_i}{m^{r+1-i}}.$$

Hence, it is sufficient to show that

$$c_1\left(R^1 f_*\left(\mathrm{Sym}^m(F \otimes f^*(H)) \otimes A^{\otimes -1} \otimes f^*(H)\right)\right)$$

is semi-ample over $S$. This can be proved in the same way as in the proof of [11, Theorem 2.4] by using [11, Proposition 2.2], Claim 2.2.2 and Proposition 1.2.1. □

Let $C$ be a smooth projective curve and $E$ a vector bundle on $C$. We say $E$ is *strongly semistable* if, for any finite morphisms $\phi : C' \to C$ of smooth projective curves, $\phi^*(E)$ is semistable. Note that if $\mathrm{char}(k) = 0$ and $E$ is semistable, then $E$ is strongly semistable. As a corollary, we have the following, which can be proved in the exactly same way as [11, Corollary 2.5].

**Corollary 2.3** ($\mathrm{char}(k) \geq 0$)**.** *Let $X$ be a quasi-projective variety, $Y$ a normal quasi-projective variety, and $f : X \to Y$ a surjective and projective morphism of* $\dim f = 1$. *Let $E$ be a locally free sheaf on $X$ and $S$ a finite subset of $Y$. If, for all $s \in S$, $f$ is flat over $s$, the geometric fiber $X_{\bar{s}}$ over $s$ is reduced and Gorenstein, and $E$ is strongly semistable on each connected component of the normalization of $X_{\bar{s}}$, then* $\mathrm{dis}_{X/Y}(E)$ *is weakly positive over $S$.*

**Remark 2.4** ($\mathrm{char}(k) = 0$)**.** In [11], we proved that the divisor

$$(8g+4)\lambda - g\delta_{irr} - \sum_{i=1}^{[g/2]} 4i(g-i)\delta_i$$

on $\bar{M}_g$ is weakly positive over any finite subsets of $M_g$. Here we give an alternative proof of this inequality.

Fix a polynomial $P_g(m) = (6m-1)(g-1)$. Let $H_g \subset \mathrm{Hilb}^{P_g}_{\mathbb{P}^{5g-6}}$ be a subscheme of all tri-canonically embedded stable curves, $Z_g \subset H_g \times \mathbb{P}^{5g-6}$ the universal tri-canonically embedded stable curves, and $f_g : Z_g \to H_g$ the natural projection. Then, $G = \mathrm{PGL}(5g-5)$ acts on $Z_g$ and $H_g$, and $f_g$ is a $G$-morphism. Let $\phi : H_g \to \bar{M}_g$ be the natural morphism of the geometric quotient. Then, by Seshadri's theorem [13, Theorem 6.1], there is a finite morphism $h : Y \to \bar{M}_g$ of normal varieties with the following properties. Let $W_g$ be the normalization of $H_g \times_{\bar{M}_g} Y$, and let $\pi : W_g \to H_g$ and $\phi' : W_g \to Y$ be the induced morphisms by the projections of $H_g \times_{\bar{M}_g} Y \to H_g$ and $H_g \times_{\bar{M}_g} Y \to Y$ respectively. Then, we have the following.

(1) $G$ acts on $W_g$, and $\pi$ is a $G$-morphism.
(2) $\phi' : W_g \to Y$ is a principal $G$-bundle.

Thus, $f'_g : U_g = Z_g \times_{H_g} W_g \to W_g$ is a stable curve, $G$ acts on $U_g$ and $f'_g$ is a $G$-morphism. Since $\phi' : W_g \to Y$ is a principal $G$-bundle, we can easily see that $U_g$ is also a principal $G$-bundle and the geometric quotient $X = U_g/G$ gives rise to a stable curve $f : X \to Y$ over $Y$. Moreover, $U_g = W_g \times_Y X$. Then, we have the following commutative diagram:

$$\begin{tikzcd}
 & Z_g \arrow[dl, "f_g"'] & & U_g \arrow[ll, "\pi'"'] \arrow[dl, "f'_g"'] \arrow[dd, "\phi''"] \\
H_g \arrow[dd, "\phi"'] & & W_g \arrow[ll, "\pi"] \arrow[dd, "\phi'"'] & \\
 & & & X \arrow[dl, "f"] \\
\bar{M}_g & & Y \arrow[ll, "h"] &
\end{tikzcd}$$

Let $\Delta$ be the minimal closed subset of $H_g$ such that $f_g$ is not smooth over a point of $\Delta$. Then, by [2, Theorem (1.6) and Corollary (1.9)], $Z_g$ and $H_g$ are quasi-projective and smooth, and $\Delta$ is a divisor with only normal crossings. Let $\Delta = \Delta_{irr} \cup \Delta_1 \cup \cdots \cup \Delta_{[g/2]}$ be the irreducible decomposition of $\Delta$ such that, if $x \in \Delta_i \setminus \operatorname{Sing}(\Delta)$ (resp. $x \in \Delta_{irr} \setminus \operatorname{Sing}(\Delta)$), then $f_g^{-1}(x)$ is a stable curve with one node of type $i$ (resp. irreducible stable curve with one node).

Form now on, we consider everything over $\bar{M}_g^{[1]}$. (Recall that $\bar{M}_g^{[1]}$ is the set of stable curves with one node at most.) In the following, the superscript "0" means the objects over $\bar{M}_g^{[1]}$.

In [10, §3], we constructed a locally sheaf $F$ on $Z_g^0$ with the following properties.

(a) $F$ is invariant by the action of $G$.

(b) For each $y \in H_g^0 \setminus (\Delta_1 \cup \cdots \cup \Delta_{[g/2]})$, $F|_{f_g^{-1}(y)} = \operatorname{Ker}\left(H^0(\omega_{f_g^{-1}(y)}) \otimes \mathcal{O}_{f_g^{-1}(y)} \to \omega_{f_g^{-1}(y)}\right)$, which is semistable on $f_g^{-1}(y)$.

(c) $\operatorname{dis}_{Z_g^0/H_g^0}(F) = (8g+4)\det(\pi_*(\omega_{Z_g^0/H_g^0})) - g\Delta_{irr}^0 - \sum_{i=1}^{[\frac{g}{2}]} 4i(g-i)\Delta_i^0$.

Then, $\pi'^*(F)$ is a $G$-invariant locally free sheaf on $U_g^0$, so that $\pi'^*(F)$ can be descended to $X^0$ because $U_g \to X$ is a principal $G$-bundle. Namely, there is a locally free sheaf $F'$ on $X^0$ such that $\phi''^*(F') = \pi'^*(F)$. Therefore, by Lemma 1.3.1, $\phi'^*(\operatorname{dis}_{X^0/Y^0}(F')) = \pi^*(\operatorname{dis}_{Z_g^0/H_g^0}(F))$. On the other hand, if we set

$$D = (8g+4)\lambda - g\delta_{irr} - \sum_{i=1}^{[\frac{g}{2}]} 4i(g-i)\delta_i,$$

then $\phi^*(D^0) = \operatorname{dis}_{Z_g^0/H_g^0}(F)$. Therefore, we get $\phi'^*(h^*(D^0)) = \phi'^*(\operatorname{dis}_{X^0/Y^0}(F'))$, which implies that $h^*(D^0) = \operatorname{dis}_{X^0/Y^0}(F')$ because $\operatorname{Pic}(W_g)^G = \operatorname{Pic}(Y)$. Moreover, by Corollary 2.3, $\operatorname{dis}_{X^0/Y^0}(F')$ is weakly positive over any finite subsets of $h^{-1}(M_g)$. Thus, $h_*(\operatorname{dis}_{X^0/Y^0}(F')) = \deg(h)D^0$ is weakly positive over any finite subsets of $M_g$ by (2) of Proposition 1.1.2. Hence, $D$ is weakly positive over any finite subsets of of $M_g$ because $\operatorname{codim}(\bar{M}_g \setminus \bar{M}_g^{[1]}) \geq 2$.

## 3. A CERTAIN KIND OF HYPERELLIPTIC FIBRATIONS

We say $f : X \to Y$ is a hyperelliptic fibered surface of genus $g$ if $X$ is a smooth projective surface, $Y$ is a smooth projective curve, the generic fiber of $f$ is a smooth hyperelliptic curve of genus

$g$, and there is no $(-1)$-curve along each fiber of $f$. Since $f$ is minimal, the hyperelliptic involution of the generic fiber extends to an automorphism of $X$ over $Y$. We denote this automorphism by $j$. Clearly, the order of $j$ is 2, namely, $j \neq \mathrm{id}_X$ and $j^2 = \mathrm{id}_X$.

The purpose of this section is to show the existence of a special kind of hyperelliptic fibered surfaces as described in the following propositions.

**Proposition 3.1** ($\mathrm{char}(k) = 0$)**.** *For fixed integers $g$ and $i$ with $g \geq 2$ and $0 \leq i \leq g-1$, there is a hyperelliptic fibered surface $f : X \to Y$ of genus $g$, and a section $\Gamma$ of $f$ such that*

1. $\mathrm{Sing}(f) \neq \emptyset$, $j(\Gamma) = \Gamma$,
2. *every singular fiber of $f$ is a stable curve consisting a smooth projective curve of genus $i$ and a smooth projective curve of genus $g-i$ meeting transversally at one point, and that*
3. *$\Gamma$ intersects with the component of genus $g-i$ on every singular fiber.*

**Proposition 3.2** ($\mathrm{char}(k) = 0$)**.** *For fixed integers $g$ and $i$ with $g \geq 2$ and $0 \leq i \leq g$, there is a hyperelliptic fibered surface $f : X \to Y$ of genus $g$, and a section $\Gamma$ of $f$ such that*

1. $\mathrm{Sing}(f) \neq \emptyset$, $j(\Gamma) \cap \Gamma = \emptyset$,
2. *every singular fiber of $f$ is a stable curve consisting a smooth projective curve of genus $i$ and a smooth projective curve of genus $g-i$ meeting transversally at one point, and that*
3. *$\Gamma$ intersects with the component of genus $g-i$ on every singular fiber.*

**Proposition 3.3** ($\mathrm{char}(k) = 0$)**.** *For fixed integers $g$ and $i$ with $g \geq 2$ and $0 \leq i \leq g-1$, there is a hyperelliptic fibered surface $f : X \to Y$ of genus $g$, and a section $\Gamma$ of $f$ such that*

1. $\mathrm{Sing}(f) \neq \emptyset$, $j(\Gamma) = \Gamma$,
2. *every singular fiber of $f$ is a semi-stable curve consisting a smooth projective curve of genus $i$ and a smooth projective curve of genus $g-i-1$ meeting transversally at two points, and that*
3. *$\Gamma$ intersects with the component of genus $g-i-1$ on every singular fiber.*

**Proposition 3.4** ($\mathrm{char}(k) = 0$)**.** *For fixed integers $g$ and $i$ with $g \geq 2$ and $0 \leq i \leq g-1$, there is a hyperelliptic fibered surface $f : X \to Y$ of genus $g$, and a section $\Gamma$ of $f$ such that*

1. $\mathrm{Sing}(f) \neq \emptyset$, $j(\Gamma) \cap \Gamma = \emptyset$,
2. *every singular fiber of $f$ is a semi-stable curve consisting a smooth projective curve of genus $i$ and a smooth projective curve of genus $g-i-1$ meeting transversally at two points, and that*
3. *$\Gamma$ intersects with the component of genus $g-i-1$ on every singular fiber.*

**Proposition 3.5** ($\mathrm{char}(k) = 0$)**.** *For fixed integers $g$ and $i$ with $g \geq 2$ and $1 \leq i \leq g-1$, there is a hyperelliptic fibered surface $f : X \to Y$ of genus $g$, and non-crossing sections $\Gamma_1$ and $\Gamma_2$ of $f$ such that*

1. $\mathrm{Sing}(f) \neq \emptyset$, $j(\Gamma_1) = \Gamma_1$, $j(\Gamma_2) = \Gamma_2$,
2. *every singular fiber of $f$ is a stable curve consisting a smooth projective curve of genus $i$ and a smooth projective curve of genus $g-i$ meeting transversally at one point,*
3. *$\Gamma_1$ and $\Gamma_2$ gives rise to a 2-pointed stable curve $(f : X \to Y, \Gamma_1, \Gamma_2)$, and that*
4. *the type of $x$ in $f^{-1}(f(x))$ as 2-pointed stable curve is $\{(i, \{1\}), (g-i, \{2\})\}$ for all $x \in \mathrm{Sing}(f)$.*

Let us begin with the following lemma.

**Lemma 3.6** ($\mathrm{char}(k) = 0$)**.** *For non-negative integers $a_1$ and $a_2$, there are a morphism $f_1 : X_1 \to Y_1$ of smooth projective varieties, an effective divisor $D_1$ on $X_1$, a line bundle $L_1$ on $X_1$, a line bundle $M_1$ on $Y_1$, and non-crossing sections $\Gamma_1$ and $\Gamma_2$ of $f_1 : X_1 \to Y_1$ with the following properties.*

(1) $\dim X_1 = 2$ *and* $\dim Y_1 = 1$.

(2) *Let $\Sigma_1$ be the set of all critical values of $f_1$, i.e., $P \in \Sigma_1$ if and only if $f_1^{-1}(P)$ is a singular variety. Then, for any $P \in Y_1 \setminus \Sigma_1$, $f_1^{-1}(P)$ is a smooth rational curve.*

(3) *$\Sigma_1 \neq \emptyset$, and for any $P \in \Sigma_1$, $f_1^{-1}(P)$ is a reduced curve consisting of two smooth rational curves $E_P^1$ and $E_P^2$ joined at one point transversally.*

(4) *$D_1$ is smooth and $f_1|_{D_1} : D_1 \to Y_1$ is etale.*

(5) $(E_P^1 \cdot D_1) = a_1 + 1$ *and* $(E_P^2 \cdot D_1) = a_2 + 1$ *for any $P \in \Sigma_1$. Moreover, $D_1$ does not pass through $E_P^1 \cap E_P^2$.*

(6) *There is a map $\sigma : \Sigma_1 \to \{1, 2\}$ such that*

$$D_1 \in \left| L_1^{\otimes a_1+a_2+2} \otimes f_1^*(M_1) \otimes \mathcal{O}_{X_1}\left( - \sum_{P\in\Sigma_1} (a_{\sigma(P)} + 1) E_P^{\sigma(P)} \right) \right|.$$

(7) $\deg(M_1)$ *is divisible by* $(a_1 + 1)(a_2 + 1)$.

(8)

$$\Gamma_1 \in \left| L_1 \otimes \mathcal{O}_{X_1}\left( - \sum_{\substack{P\in\Sigma_1 \\ \sigma(P)=1}} E_P^1 \right) \right| \quad \textit{and} \quad \Gamma_2 \in \left| L_1 \otimes \mathcal{O}_{X_1}\left( - \sum_{\substack{P\in\Sigma_1 \\ \sigma(P)=2}} E_P^2 \right) \right|.$$

*Moreover,*

$$(D_1 \cdot \Gamma_1) = (D_1 \cdot \Gamma_2) = 0 \quad \textit{and} \quad (E_P^i \cdot \Gamma_j) = \begin{cases} 0 & \textit{if } i \neq j, \\ 1 & \textit{if } i = j. \end{cases}$$

*Proof.* We can prove this lemma in the exactly same way as in [11, Lemma A.1] with a slight effort. We use the notation in [11, Lemma A.1]. Let $F_1$ and $F_2$ be curves in $\mathbb{P}^1_{(X,Y)} \times \mathbb{P}^1_{(S,T)}$ defined by $\{X = 0\}$ and $\{X = Y\}$ respectively. Note that $F_1 = p^{-1}((0 : 1))$, $F_2 = p^{-1}((1 : 1))$, $D'' = p^{-1}((1 : 0))$, $(D' \cdot F_1) = (D' \cdot F_2) = 1$, $D' \cap F_1 = \{Q_1\}$ and $D' \cap F_2 = \{Q_2\}$. Then, since $u_1^*(F_1) = p_1^{-1}((0 : 1))$ and $u_1^*(F_2) = p_1^{-1}((1 : 1))$, in [11, Claim A.1.3], we can see that each tangent of $u_1^*(D)$ at $Q_{i,j}$ ($i = 1, 2$) is different from $u_1^*(F_i)$.

Let $\overline{\Gamma}_i$ be the strict transform of $u_1^*(F_i)$ by $\mu_1 : Z_1 \to \mathbb{P}^1_{(X,Y)} \times Y$. Then,

$$\overline{\Gamma}_i \in \left| \mu_1^*(p_1^*(\mathcal{O}_{\mathbb{P}^1}(1))) \otimes \mathcal{O}_{Z_1}\left( - \sum_j E_{i,j} \right) \right|.$$

Thus, if we set $\Gamma_i = (\nu_1)_*(\overline{\Gamma}_i)$, then we get our lemma. □

In the following proofs, we use the notation in [11, Proposition A.2 and Proposition A.3].

**The proof of Proposition 3.1:** We apply Lemma 3.6 to the case where $a_1 = 2i$ and $a_2 = 2g - 2i - 1$. We replace $D_2$ by $D_2 + \Gamma_2$ and $a_2$ by $a_2 + 1$. Then, (4), (5) and (6) hold for the new

$D_2$ and $a_2$. Thus, we can construct $f : X \to Y$ in exactly same way as in [11, Proposition A.2]. Since $u_2^*(\Gamma_2)$ is the ramification locus of $\mu_3$, $\overline{\Gamma} = h^*(u_2^*(\Gamma_2))_{red}$ is a section of $f_3$. Thus, if we set $\Gamma = \nu_3(\overline{\Gamma})$, then we have our desired example.

**The proof of Proposition 3.2:** Applying Lemma 3.6 to the case where $a_1 = 2i$ and $a_2 = 2g-2i$, we can construct $f : X \to Y$ in exactly same way as in [11, Proposition A.2]. Here let us consider $u_2^*(\Gamma_2)$. Then, $u_2^*(\Gamma_2)$ is a section of $f_2$ such that $u_2^*(\Gamma_2) \cap (D_2 + B) = \emptyset$, $(u_2^*(\Gamma_2) \cdot \overline{E}_Q^1) = 0$ and $(u_2^*(\Gamma_2) \cdot \overline{E}_Q^2) = 1$ for all $Q \in \Sigma_2$. Here we set $\Gamma' = \nu_3(\mu_3^*(u_2^*(\Gamma_2)))$. Then, since $\mu_3^*(u_2^*(\Gamma_2))$ does not intersect with the ramification locus of $\mu_3$, $\Gamma'$ is etale over $Y$. Moreover, we can see $(\Gamma' \cdot C_Q^1) = 0$ and $(\Gamma' \cdot C_Q^2) = 2$ for all $Q \in \Sigma_2$. If $\Gamma'$ is not irreducible, then we choose $\Gamma$ as one of irreducible component of $\Gamma'$. If $\Gamma'$ is irreducible, then we consider $X \times_Y \Gamma \to \Gamma$ and the natural section of $X \times_Y \Gamma \to \Gamma$. Then we get our desired example.

**The proof of Proposition 3.3:** We apply Lemma 3.6 to the case where $a_1 = 2i + 1$ and $a_2 = 2g - 2i - 2$. We replace $D_2$ by $D_2 + \Gamma_2$ and $a_2$ by $a_2 + 1$. Then, (4), (5) and (6) hold for the new $D_2$ and $a_2$. Note that $\deg(M_1)$ is even. Thus, we can get a double covering $\mu : X \to X_1$ in exactly same way as in [11, Proposition A.3]. Let $f : X \to Y_1$ be the induced morphism, and $\Gamma = \mu^*(\Gamma_2)_{red}$. Then, we have our desired example.

**The proof of Proposition 3.4:** Applying Lemma 3.6 to the case where $a_1 = 2i+1$ and $a_2 = 2g-2i-1$, we can get a double covering $\mu : X \to X_1$ in exactly same way as in [11, Proposition A.3]. Let $f : X \to Y_1$ be the induced morphism and $\Gamma' = \mu^*(\Gamma_2)$. Then, $\Gamma'$ is etale over $Y_1$. If $\Gamma'$ is not irreducible, then we choose $\Gamma$ as one of irreducible component of $\Gamma'$. If $\Gamma'$ is irreducible, then we consider $X \times_{Y_1} \Gamma \to \Gamma$ and the natural section of $X \times_{Y_1} \Gamma \to \Gamma$. Then we get our desired example.

**The proof of Proposition 3.5:** We apply Lemma 3.6 to the case where $a_1 = 2i - 1$ and $a_2 = 2g - 2i - 1$. We replace $D_1$ by $D_1 + \Gamma_1$, $D_2$ by $D_2 + \Gamma_2$, $a_1$ by $a_1 + 1$, and $a_2$ by $a_2 + 1$. Then, (4), (5) and (6) hold for the new $D_1$, $D_2$, $a_1$ and $a_2$. Thus, we can construct $f : X \to Y$ in exactly same way as in [11, Proposition A.2]. Since $u_2^*(\Gamma_1)$ and $u_2^*(\Gamma_2)$ are the ramification locus of $\mu_3$, $\overline{\Gamma}_1 = h^*(u_2^*(\Gamma_1))_{red}$ and $\overline{\Gamma}_2 = h^*(u_2^*(\Gamma_2))_{red}$ are sections of $f_3$. Thus, if we set $\Gamma_1 = \nu_3(\overline{\Gamma}_1)$ and $\Gamma_2 = \nu_3(\overline{\Gamma}_2)$, then we have our desired example.

Finally, let us consider the following two lemmas, which will be used in the later section.

**Lemma 3.7** ($\mathrm{char}(k) \geq 0$). *Let $X$ be a smooth projective surface and $Y$ a smooth projective curve. Let $f : X \to Y$ be a surjective morphism with connected fibers, and let $L$ be a line bundle on $X$. If $L|_{X_\eta}$ gives rise to a torsion element of $\mathrm{Pic}(X_\eta)$ on the generic fiber $X_\eta$ of $f$ and $\deg(L|_F) = 0$ for every irreducible component $F$ of fibers, then we have $(L^2) = 0$.*

*Proof.* Replacing $L$ by $L^{\otimes n}$ ($n \neq 0$), we may assume that $L|_{X_\eta} \simeq \mathcal{O}_{X_\eta}$. Thus, $f_*(L)$ is a line bundle on $Y$, and the natural homomorphism $f^*f_*(L) \to L$ is injective. Hence, there is an effective divisor $E$ on $X$ such that $f^*f_*(L) \otimes \mathcal{O}_X(E) \simeq L$. Since $f^*f_*(L) \to L$ is surjective on the generic fiber, $E$ is a vertical divisor. Moreover, $(E \cdot F) = 0$ for every irreducible component $F$ of fibers. Therefore, by Zariski's lemma, $(E^2) = 0$. Hence, $(L^2) = (E^2) = 0$. □

**Lemma 3.8** ($\mathrm{char}(k) \geq 0$). *Let $C$ be a smooth projective curve of genus $g \geq 2$. Let $\vartheta$ be a line bundle on $C$ with $\vartheta^{\otimes 2} = \omega_C$. Let $\Delta$ be the diagonal of $C \times C$, and $p : C \times C \to C$ and*

*$q : C \times C \to C$ the projection to the first factor and the projection to the second factor respectively. Then, $p^*(\vartheta^{\otimes n}) \otimes q^*(\vartheta^{\otimes n}) \otimes \mathcal{O}_{C\times C}((n-1)\Delta)$ is generated by global sections for all $n \geq 3$.*

*Proof.* Since $p^*(\vartheta^{\otimes n}) \otimes q^*(\vartheta^{\otimes n})$ is generated by global sections, the base locus of $p^*(\vartheta^{\otimes n}) \otimes q^*(\vartheta^{\otimes n}) \otimes \mathcal{O}_{C\times C}((n-1)\Delta)$ is contained in $\Delta$. Moreover,

$$p^*(\vartheta^{\otimes n}) \otimes q^*(\vartheta^{\otimes n}) \otimes \mathcal{O}_{C\times C}((n-1)\Delta)\big|_{\Delta} \simeq \omega_C.$$

Thus, it is sufficient to see that

$$H^0(p^*(\vartheta^{\otimes n}) \otimes q^*(\vartheta^{\otimes n}) \otimes \mathcal{O}_{C\times C}((n-1)\Delta)) \to H^0(p^*(\vartheta^{\otimes n}) \otimes q^*(\vartheta^{\otimes n}) \otimes \mathcal{O}_{C\times C}((n-1)\Delta)\big|_{\Delta})$$

is surjective.

We define $L_{n,i}$ to be

$$L_{n,i} = p^*(\vartheta^{\otimes n}) \otimes q^*(\vartheta^{\otimes n}) \otimes \mathcal{O}_{C\times C}(i\Delta).$$

Then, it suffices to check $H^1(L_{n,n-2}) = 0$ for the above assertion. By induction on $i$, we will see that $H^1(L_{n,i}) = 0$ for $0 \leq i \leq n-2$.

First of all, note that $H^1(\vartheta^{\otimes n}) = 0$ for $n \geq 3$. Thus,

$$H^1(p^*(\vartheta^{\otimes n}) \otimes q^*(\vartheta^{\otimes n})) = H^1(p_*(p^*(\vartheta^{\otimes n}) \otimes q^*(\vartheta^{\otimes n}))) = H^1(\vartheta^{\otimes n}) \otimes H^1(\vartheta^{\otimes n}) = 0.$$

Moreover, let us consider the exact sequence

$$0 \to L_{n,i-1} \to L_{n,i} \to L_{n,i}|_{\Delta} \to 0.$$

Here since $L_{n,i}|_{\Delta} \simeq \omega_C^{\otimes n-i}$, $H^1(L_{n,i}|_{\Delta}) = 0$ if $i \leq n-2$. Thus, by the hypothesis of induction, we can see $H^1(L_{n,i}) = 0$. □

## 4. Slope inequalities on $\bar{M}_{g,T}$

Let $g$ be a non-negative integer and $T$ a finite set with $2g-2+|T| > 0$. Recall that

$$\Upsilon_{g,T} = \{(i,I) \mid i \in \mathbb{Z}, 0 \leq i \leq g \text{ and } I \subseteq T\} \setminus (\{(0,\emptyset)\} \cup \{(0,\{t\})\}_{t\in T}),$$
$$\overline{\Upsilon}_{g,T} = \{\{(i,I),(j,J)\} \mid (i,I),(j,J) \in \Upsilon_{g,T}, i+j=g, I\cap J = \emptyset, I \cup J = T\}.$$

For a subset $L$ of $T$, let us introduce a function $\gamma_L : \Upsilon_{g,T} \times \Upsilon_{g,T} \to \mathbb{Z}$ given by

$$\gamma_L((i,I),(j,J)) = \left(\det\begin{pmatrix} i & |L\cap I| \\ j & |L \cap J|\end{pmatrix} + |L\cap I|\right)\left(\det\begin{pmatrix} i & |L\cap I| \\ j & |L \cap J|\end{pmatrix} - |L\cap J|\right).$$

Note that $\gamma_L((i,I),(j,J)) = \gamma_L((j,J),(i,I))$, so that $\gamma_L$ gives rise to a function on $\overline{\Upsilon}_{g,T}$. Further, a $\mathbb{Q}$-divisor $\theta_L$ on $\bar{M}_{g,T}$ is defined to be

$$\theta_L = 4(g-1+|L|)(g-1)\sum_{t\in L}\psi_t - 12|L|^2\lambda + |L|^2\delta_{irr} - \sum_{\upsilon\in\overline{\Upsilon}_{g,T}} 4\gamma_L(\upsilon)\delta_{\upsilon}.$$

Then, we have the following.

**Theorem 4.1** ($\operatorname{char}(k) \geq 0$)**.** *For any subset $L$ of $T$, the divisor $\theta_L$ is weakly positive over any finite subsets of $M_{g,T}$.*

*Proof.* Clearly, we may assume $T = [n]$ for some non-negative integer $n$. Let us take an $n$-pointed stable curve $f : X \to Y$ such that the induced morphism $h : Y \to \bar{M}_{g,[n]}$ is a finite and surjective morphism of normal varieties. Let $Y_0$ be the maximal Zariski open set of $Y$ over which $f$ is smooth. Let $Y \setminus Y_0 = B_1 \cup \cdots \cup B_s$ be the irreducible decomposition of $Y \setminus Y_0$. By using [3, Lemma 3.2], we can take a Zariski open set $Y_1$ with the following properties.

1. $\operatorname{codim}(Y \setminus Y_1) \geq 2$ and $Y_0 \subseteq Y_1$.
2. $Y_1$ is smooth at any points of $Y_1 \cap (Y \setminus Y_0)$.
3. $f : \operatorname{Sing}(f) \cap f^{-1}(Y_1) \to f(\operatorname{Sing}(f)) \cap Y_1$ is an isomorphism, so that for all $y \in Y_1$, the number of nodes of $f^{-1}(y)$ is one at most.
4. There is a projective birational morphism $\phi : Z_1 \to X_1 = f^{-1}(Y_1)$ such that if we set $f_1 = f \cdot \phi$, then $Z_1$ is smooth at any points of $\operatorname{Sing}(f_1) \cap f_1^{-1}(Y \setminus Y_0)$ and $f_1 : Z_1 \to Y_1$ is an $n$-pointed semi-stable curve. Moreover, $\phi$ is an isomorphism over $X_1 \setminus \operatorname{Sing}(f)$.
5. For each $l = 1, \ldots, s$, there is a $t_l$ such that $\operatorname{mult}_x(X) = t_l + 1$ for all $x \in \operatorname{Sing}(f)$ with $f(x) \in B_l \cap Y_1$.

Let $K_0$ be a subset of $\{1, \ldots, s\}$ such that $f^{-1}(x)$ is irreducible for all $x \in B_l \cap Y_1$, and let $K_1 = \{1, \ldots, s\} \setminus K_0$. For each $l \in K_1$, there is a $(g_l, I_l), (h_l, J_l) \in \Upsilon_{g,[n]}$ such that the type of $x$ is $\{(g_l, I_l), (h_l, J_l)\}$ for all $x \in \operatorname{Sing}(f)$ with $f(x) \in B_l \cap Y_1$. From now on, by abuse of notation, we denote $B_l \cap Y_1$ by $B_l$. For $l \in K_1$, $f_1^{-1}(B_l)$ has two essential components $T_l^1$ and $T_l^2$, and the components of $(-2)$-curves $E_1, \ldots, E_{t_l}$ such that $T_l^1 \to B_l$ is an $I_l$-pointed smooth curve of genus $g_l$ and $T_l^2 \to B_l$ is a $J_l$-pointed smooth curve of genus $h_l$. Moreover, the numbering of $E_1, \ldots, E_{t_l}$ is arranged as the following figure:



Let $\Gamma_1, \ldots, \Gamma_n$ be the sections of the $n$-pointed stable curve of $f : X \to Y$. By abuse of notation, the lifting of $\Gamma_a$ to $Z_1$ is also denoted by $\Gamma_a$. Here we consider a line bundle $L$ on $Z_1$ given by

$$L = \omega_{Z_1/Y_1}^{\otimes |L|} \otimes \mathcal{O}_{Z_1}\left( -(2g-2)\sum_{a \in L} \Gamma_a + \sum_{l \in K_1} \left( |L|(2g_l - 1) - (2g-2)|L \cap I_l| \right) \tilde{T}_l^1 \right),$$

where

$$\tilde{T}_l^1 = (t_l + 1)T_l^1 + \sum_{a=1}^{t_l} (t_l + 1 - a) E_a.$$

We set $E = \mathcal{O}_{X_1} \oplus L$. Then, $\operatorname{dis}_{X_1/Y_1}(E) = -(f_1)_*(c_1(L)^2)$. Here, we know the following formulae:

$$\begin{cases} f_*(c_1(\omega_{Z_1/Y_1}) \cdot \tilde{T}_l^1) = (t_l+1)(2g_l-1)B_l, \\ f_*(\tilde{T}_l^1 \cdot \tilde{T}_{l'}^1) = \begin{cases} 0 & \text{if } l \neq l', \\ -(t_l+1)B_l & \text{if } l = l', \end{cases} \\ f_*(\sum_{a\in L} \Gamma_a \cdot \tilde{T}_l^1) = (t_l+1)|L \cap I_l| B_l, \\ f_*(c_1(\omega_{Z_1/Y_1}) \cdot \Gamma_a) = -f_*(\Gamma_a \cdot \Gamma_a), \quad \text{(adjunction formula)} \\ 12\det(f_*(\omega_{Z_1/Y_1})) - \sum_{l=1}^{s}(t_l+1)B_l = f_*(c_1(\omega_{Z_1/Y_1})^2) \quad \text{(Noether's formula).} \end{cases}$$

Thus, we can see that

$$\begin{aligned} \operatorname{dis}_{Z_1/Y_1}(E) = 4(g-1+|L|)(g-1)f_*\left(c_1(\omega_{Z_1/Y_1}) \cdot \sum_{a\in L}\Gamma_a\right) - 12|L|^2\det(f_*(\omega_{Z_1/Y_1})) \\ + \sum_{l\in K_0}|L|^2(t_l+1)B_l - \sum_{l\in K_1} 4(t_l+1)\gamma_L(\{(g_l,I_l),(h_l,J_l)\})B_l. \end{aligned}$$

On the other hand, for $y \in Y_0$, let $\phi : C' \to f^{-1}(y)$ be a finite morphism of smooth projective curves. Then, $\phi^*(E|_{f^{-1}(y)}) = \mathcal{O}_{C'} \oplus \phi^*(L|_{f^{-1}(y)})$ and

$$\deg(\phi^*(L|_{f^{-1}(y)})) = \deg(\phi)\deg(L|_{f^{-1}(y)}) = 0.$$

Therefore, $\phi^*(E|_{f^{-1}(y)})$ is semistable, which means that $E|_{f^{-1}(y)}$ is strongly semistable for all $y \in Y_0$. Thus, by Corollary 2.3, $\operatorname{dis}_{Z_1/Y_1}(E)$ is weakly positive over any finite subsets of $Y_0$ as a divisor on $Y_1$. Therefore, if we set

$$\begin{aligned} \theta'_L = 4(g-1+|L|)(g-1)f_*\left(c_1(\omega_{Z_1/Y_1}) \cdot \sum_{a\in L}\Gamma_a\right) - 12|L|^2\det(f_*(\omega_{Z_1/Y_1})) \\ + \sum_{l\in K_0}|L|^2(t_l+1)B_l - \sum_{l\in K_1} 4(t_l+1)\gamma_L(\{(g_l,I_l),(h_l,J_l)\})B_l. \end{aligned}$$

on $Y$, then $\theta'_L$ is weakly positive over any finite subsets of $Y_0$ as a divisor on $Y$. Here $h^*(\theta_L) = \theta'_L$, so that $h_*(\theta'_L) = \deg(h)\theta_L$ by the projection formula. Hence we have our theorem by (2) of Proposition 1.1.2. □

Let us apply Theorem 4.1 to the cases $\bar{M}_{g,1}$ and $\bar{M}_{g,2}$.

**Corollary 4.2** ($\operatorname{char}(k) = 0$)**.** *Let $\bar{M}_{g,1} = \bar{M}_{g,\{1\}}$ be the moduli space of one-pointed stable curves of genus $g \geq 1$. We set $\delta_i, \mu, \theta_1 \in \operatorname{Pic}(\bar{M}_{g,1}) \otimes \mathbb{Q}$ as follows:*

$$\begin{cases} \delta_i = \delta_{\{(i,\emptyset),(g-i,\{1\})\}} \quad (1 \leq i \leq g-1) \\ \mu = (8g+4)\lambda - g\delta_{irr} - \sum_{i=1}^{g-1} 4i(g-i)\delta_i \\ \theta_1 = 4g(g-1)\psi_1 - 12\lambda + \delta_{irr} - \sum_{i=1}^{g-1} 4i(i-1)\delta_i. \end{cases}$$

*Then, we have the following:*

(1) *$\mu$ and $\theta_1$ are weakly positive over any finite subsets of $M_{g,1}$. In particular,*

$$\mathbb{Q}_+\mu + \mathbb{Q}_+\theta_1 + \mathbb{Q}_+\delta_{irr} + \sum_{i=1}^{g-1} \mathbb{Q}_+\delta_i \subseteq \operatorname{Nef}(\bar{M}_{g,1}; M_{g,1}),$$

*where $\mathbb{Q}_+ = \{x \in \mathbb{Q} \mid x \geq 0\}$. (Note that $\mu = \theta_1 = 0$ if $g = 1$, and $\mu = 0$ if $g = 2$.)*

(2) *We assume $g = 1$. Then, $a\mu + b\theta_1 + c_{irr}\delta_{irr}$ is nef over $M_{1,1}$ if and only if $c_{irr} \geq 0$.*

(3) *We assume $g \geq 2$. If a $\mathbb{Q}$-divisor*

$$D = a\mu + b\theta_1 + c_{irr}\delta_{irr} + \sum_{i=1}^{g-1} c_i\delta_i$$

*is nef over $M_{g,1}$, then $b, c_{irr}, c_1, \dots, c_{g-1}$ are non-negative.*

*Proof.* (1) $\mu$ is weakly positive over any finite subsets of $M_{g,1}$ by [11, Theorem B] or Remark 2.4, and (2) of Proposition 1.1.3. Moreover, $\theta_1$ is weakly positive over any finite subsets of $M_{g,1}$ by virtue of the case $T = L = \{1\}$ in Theorem 4.1.

(2) This is obvious because $\mu = \theta_1 = 0$.

(3) We assume that $D$ is nef over $M_{g,1}$. Let $C$ be a smooth curve of genus $g$, and $\Delta$ the diagonal of $C \times C$. Let $p : C \times C \to C$ be the projection to the first factor. Then, $\Delta$ gives rise to a section of $p$. Hence, we get a morphism $\varphi_1 : C \to \bar{M}_{g,1}$ with $\varphi_1(C) \subseteq M_{g,1}$. By our assumption, $\deg(\varphi_1^*(D)) \geq 0$. On the other hand,

$$\deg(\varphi_1^*(\mu)) = \deg(\varphi_1^*(\delta_{irr})) = \deg(\varphi_1^*(\delta_1)) = \cdots = \deg(\varphi_1^*(\delta_{g-1})) = 0$$

and $\deg(\varphi_1^*(\theta_1)) = 8g(g-1)^2$. Thus, $b \geq 0$.

Let $f_2 : X_2 \to Y_2$ be a hyperelliptic fibered surface and $\Gamma_2$ a section as in Proposition 3.3 for $i = 0$. Let $\varphi_2 : Y_2 \to \bar{M}_{g,1}$ be the induced morphism. Then, $\varphi_2(Y_2) \cap M_{g,1} \neq \emptyset$,

$$\deg(\varphi_2^*(\mu)) = \deg(\varphi_2^*(\delta_1)) = \cdots = \deg(\varphi_2^*(\delta_{g-1})) = 0$$

and $\deg(\varphi_2^*(\delta_{irr})) = \deg(\delta_{irr}(X_2/Y_2)) > 0$. On the generic fiber, $\Gamma_2$ is a ramification point of the hyperelliptic covering. Thus,

$$L_2 = \omega_{X_2/Y_2} \otimes \mathcal{O}_{X_2}(-(2g-2)\Gamma_2)$$

satisfies the conditions of Lemma 3.7. Thus, $(L_2^2) = 0$, which says us that $\deg(\varphi_2^*(\theta_1)) = 0$. Therefore, we get $c_{irr} \geq 0$.

Finally we fix $i$ with $1 \leq i \leq g-1$. Let $f_3 : X_3 \to Y_3$ be a hyperelliptic fibered surface and $\Gamma_3$ a section as in Proposition 3.1. Let $\varphi_3 : Y_3 \to \bar{M}_{g,1}$ be the induced morphism. Then, $\varphi_3(Y_3) \cap M_{g,1} \neq \emptyset$, $\deg(\varphi_3^*(\mu)) = 0$, $\deg(\varphi_3^*(\delta_l)) = 0$ ($l \neq i$) and $\deg(\varphi_3^*(\delta_i)) = \deg(\delta_i(X_3/Y_3)) > 0$. Let $\Sigma_3$ be the set of critical values of $f_3$. For each $P \in \Sigma_3$, let $E_P$ be the component of genus $i$ in $f_3^{-1}(P)$. On the generic fiber, $\Gamma_2$ is a ramification point of the hyperelliptic covering. Thus,

$$L_3 = \omega_{X_3/Y_3} \otimes \mathcal{O}_{X_3}\left(-(2g-2)\Gamma_3 + \sum_{P \in \Sigma_3} (2i-1)E_P\right)$$

satisfies the conditions of Lemma 3.7. Therefore, $(L_3^2) = 0$, which says us that $\deg(\varphi_3^*(\theta_1)) = 0$. Hence, we get $c_i \geq 0$. □

**Corollary 4.3** ($\operatorname{char}(k) = 0$)**.** *Let $\bar{M}_{g,2} = \bar{M}_{g,\{1,2\}}$ be the moduli space of two-pointed stable curves of genus $g \geq 2$. We set $\delta_i, \sigma_i, \mu, \theta_{1,2} \in \operatorname{Pic}(\bar{M}_{g,2}) \otimes \mathbb{Q}$ as follows:*

$$\begin{cases} \delta_i = \delta_{\{(i,\emptyset),(g-i,\{1,2\})\}} \quad (1 \leq i \leq g) \\ \sigma_i = \delta_{\{(i,\{1\}),(g-i,\{2\})\}} \quad (1 \leq i \leq g-1) \\ \mu = (8g+4)\lambda - g\delta_{irr} - \sum_{i=1}^{g-1} 4i(g-i)\sigma_i - \sum_{i=1}^{g} 4i(g-i)\delta_i, \\ \theta_{1,2} = (g-1)(g+1)(\psi_1+\psi_2) - 12\lambda + \delta_{irr} \\ \qquad\qquad - \sum_{i=1}^{g-1}(2i-g-1)(2i-g+1)\sigma_i - \sum_{i=1}^{g} 4i(i-1)\delta_i. \end{cases}$$

*Then, we have the following.*

(1) *$\mu$ and $\theta_{1,2}$ are weakly positive over any finite subsets of $M_{g,2}$. In particular,*

$$\mathbb{Q}_+\mu + \mathbb{Q}_+\theta_{1,2} + \mathbb{Q}_+\delta_{irr} + \sum_{i=1}^{g-1} \mathbb{Q}_+\sigma_i + \sum_{i=1}^{g} \mathbb{Q}_+\delta_i \subseteq \operatorname{Nef}(\bar{M}_{g,2}; M_{g,2}).$$

(2) *If a $\mathbb{Q}$-divisor*

$$D = a\mu + b\theta_{1,2} + c_{irr}\delta_{irr} + \sum_{i=1}^{g-1} c_i\sigma_i + \sum_{i=1}^{g} d_i\delta_i$$

*on $\bar{M}_{g,2}$ is nef over $M_{g,2}$, then*

$$b \geq 0,\ c_{irr} \geq 0,\ c_i \geq 0\ (\forall i = 1, \dots, g-1),\ d_i \geq 0\ (\forall i = 1, \dots, g).$$

(3) *Here we set $\sigma$, $\mu'$ and $\theta'_{1,2}$ as follows:*

$$\begin{cases} \sigma = \delta_{irr} + \sum_{i=1}^{g-1} \sigma_i, \\ \mu' = (8g+4)\lambda - g\sigma - \sum_{i=1}^{g} 4i(g-i)\delta_i, \\ \theta'_{1,2} = (g-1)(g+1)(\psi_1+\psi_2) - 12\lambda + \sigma - \sum_{i=1}^{g} 4i(i-1)\delta_i. \end{cases}$$

*Then, we have*

$$\mathbb{Q}_+\mu' + \mathbb{Q}_+\theta'_{1,2} + \mathbb{Q}_+\sigma + \sum_{i=1}^{g} \mathbb{Q}_+\delta_i \subseteq \operatorname{Nef}(\bar{M}_{g,2}; M_{g,2}).$$

*Moreover, if a $\mathbb{Q}$-divisor $a\mu' + b\theta'_{1,2} + c\sigma + \sum_{i=1}^{g} d_i\delta_i$ on $\bar{M}_{g,2}$ is nef over $M_{g,2}$, then $b$, $c$, $d_1, \dots, d_g$ are non-negative.*

*Proof.* (1) By [11, Theorem B] or Remark 2.4, and (2) of Proposition 1.1.3, $\mu$ is weakly positive over any finite subsets of $M_{g,2}$. Further, $\theta_{1,2}$ is weakly positive over any finite subsets of $M_{g,2}$ by the case $T = L = \{1, 2\}$ in Theorem 4.1.

(2) We assume that $D$ is nef over $M_{g,2}$. Let $C$ be a smooth curve of genus $g$, and $\Delta$ the diagonal of $C \times C$. Let $p : C \times C \to C$ and $q : C \times C \to C$ be the projection to the first factor and the second factor respectively. Moreover, let $\vartheta$ be a line bundle on $C$ with $\vartheta^{\otimes 2} = \omega_C$ and $L_n = p^*(\vartheta^{\otimes n}) \otimes q^*(\vartheta^{\otimes n}) \otimes \mathcal{O}_{C\times C}((n-1)\Delta)$. For $n \geq 3$, let $T_n$ be a general member of $|L_n|$. Then, since $(L_n^2) > 0$, by Lemma 3.8, $T_n$ is smooth and irreducible. Moreover, $T_n$ meets $\Delta$ transversally. Then, we have two morphisms $p_n : T_n \to C$ and $q_n : T_n \to C$ given by $T_n \hookrightarrow C \times C \xrightarrow{p} C$ and $T_n \hookrightarrow C \times C \xrightarrow{q} C$ respectively. Let $\Gamma_{p_n}$ and $\Gamma_{q_n}$ be the graph of $p_n$ and $q_n$ in $C \times T_n$ respectively.

Then, it is easy to see that $\Gamma_{p_n}$ and $\Gamma_{q_n}$ meet transversally, and $(\Gamma_{p_n} \cdot \Gamma_{q_n}) = (T_n \cdot \Delta) = 2g-2$. Let $X \to C \times T_n$ be the blowing-ups at points in $\Gamma_{p_n} \cap \Gamma_{q_n}$, and let $\overline{\Gamma}_{p_n}$ and $\overline{\Gamma}_{q_n}$ be the strict transform of $\Gamma_{p_n}$ and $\Gamma_{q_n}$ respectively. Then, $\overline{\Gamma}_{p_n}$ and $\overline{\Gamma}_{q_n}$ give rise to two non-crossing sections of $X \to T_n$. Moreover,

$$(\omega_{X/T_n} \cdot \overline{\Gamma}_{p_n}) = (\omega_{C \times T_n/T_n} \cdot \Gamma_{p_n}) = 2(g-1)\deg(\Gamma_{p_n} \to C) = 2(g-1)(ng-1).$$

In the same way, $(\omega_{X/T_n} \cdot \overline{\Gamma}_{q_n}) = 2(g-1)(ng-1)$. Let $\pi_n : T_n \to \bar{M}_{g,2}$ be the induced morphism. Then, we can see that $\deg(\pi_n^*(\lambda)) = \deg(\pi_n^*(\sigma_i)) = \deg(\pi_n^*(\delta_i)) = 0$ for all $i = 1, \ldots, g-1$. Moreover, $\deg(\pi_n^*(\psi_1 + \psi_2)) = 4(g-1)(ng-1)$, and $\deg(\pi_n^*(\delta_g)) = 2(g-1)$. Thus,

$$\deg(\pi_n^*(D)) = 4(g+1)(g-1)^2(ng-1)b - 8g(g-1)^2 d_g \geq 0$$

for all $n \geq 3$. Therefore, we get $b \geq 0$.

Let $f_2 : X_2 \to Y_2$ be a hyperelliptic fibered surface and $\Gamma_2$ a section as in Proposition 3.4 for $i = 0$. Then, $\Gamma_2$ and $j(\Gamma_2)$ gives two points of $X_2$ over $Y_2$. Let $\varphi_2 : Y_2 \to \bar{M}_{g,2}$ be the induced morphism. Then, $\varphi_2(Y_2) \cap M_{g,2} \neq \emptyset$, $\deg(\varphi_2^*(\mu)) = 0$, $\deg(\varphi_2^*(\sigma_i)) = 0$ for all $i = 1, \ldots, g-1$, and $\deg(\varphi_2^*(\delta_i)) = 0$ for all $i = 1, \ldots, g$. Moreover, $\deg(\varphi_2(\delta_{irr})) > 0$. On the generic fiber, two points arising from $\Gamma_2$ and $j(\Gamma_2)$ are invariant under the action of the hyperelliptic involution. Thus,

$$L_2 = \omega_{X_2/Y_2} \otimes \mathcal{O}_{X_2}(-(g-1)(\Gamma_2 + j(\Gamma_2))$$

satisfies the conditions of Lemma 3.7. Thus, $(L_2^2) = 0$, which says us that $\deg(\varphi_2^*(\theta_{1,2})) = 0$. Thus, we get $c_{irr} \geq 0$.

We fix $i$ with $1 \leq i \leq g$. Let $f_3 : X_3 \to Y_3$ be a hyperelliptic fibered surface and $\Gamma_3$ a section as in Proposition 3.2. Let $\varphi_3 : Y_3 \to \bar{M}_{g,2}$ be the induced morphism arising from the 2-pointed curve $\{f_3 : X_3 \to Y_3; \Gamma_3, j(\Gamma_3)\}$. Then, $\varphi_3(Y_3) \cap M_{g,2} \neq \emptyset$, $\deg(\varphi_3^*(\mu)) = 0$, $\deg(\varphi_3^*(\sigma_s)) = 0$ ($\forall s = 1, \ldots, g-1$), $\deg(\varphi_3^*(\delta_s)) = 0$ ($\forall s \neq i$) and $\deg(\varphi_3^*(\delta_i)) = \deg(\delta_i(X_3/Y_3)) > 0$. Let $\Sigma_3$ be the set of critical values of $f_3$. For each $P \in \Sigma_3$, let $E_P$ be the component of genus $i$ in $f_3^{-1}(P)$. On the generic fiber, two points arising from $\Gamma_2$ and $j(\Gamma_2)$ are invariant under the action of the hyperelliptic involution. Thus,

$$L_3 = \omega_{X_3/Y_3} \otimes \mathcal{O}_{X_3}\left(-(g-1)(\Gamma_3 + j(\Gamma_3)) + \sum_{P \in \Sigma_3}(2i-1)E_P\right)$$

satisfies the conditions of Lemma 3.7. Therefore, $(L_3^2) = 0$, which says us that $\deg(\varphi_3^*(\theta_{1,2})) = 0$. Hence, we get $d_i \geq 0$.

Finally we fix $i$ with $1 \leq i \leq g-1$. Let $f_4 : X_4 \to Y_4$ be a hyperelliptic fibered surface and $\Gamma_4, \Gamma_4'$ sections as in Proposition 3.5. Let $\varphi_4 : Y_4 \to \bar{M}_{g,2}$ be the induced morphism. Then, $\varphi_4(Y_4) \cap M_{g,2} \neq \emptyset$, $\deg(\varphi_4^*(\mu)) = 0$, $\deg(\varphi_4^*(\delta_s)) = 0$ ($\forall s$), $\deg(\varphi_4^*(\sigma_s)) = 0$ ($\forall s \neq i$), and $\deg(\varphi_4^*(\sigma_i)) > 0$. Let $\Sigma_4$ be the set of critical values of $f_4$. For each $P \in \Sigma_4$, let $E_P$ be the component of genus $i$ in $f_4^{-1}(P)$. On the generic fiber, $\Gamma_4$ and $\Gamma_4'$ are a ramification point of the hyperelliptic covering. Thus,

$$L_4 = \omega_{X_4/Y_4} \otimes \mathcal{O}_{X_4}\left(-(g-1)(\Gamma_4 + \Gamma_4') + \sum_{P \in \Sigma_4}\left((2i-1) - (g-1)\right)E_P\right)$$

satisfies the conditions of Lemma 3.7. Therefore, $(L_4^2) = 0$, which says us that $\deg(\varphi_4^*(\theta_{1,2})) = 0$. Hence, we get $c_i \geq 0$.

(3) There are non-negative integers $e_i$ and $f_i$ ($1 \leq i \leq g-1$) with

$$\mu' = \mu + \sum_{i=1}^{g-1} e_i \sigma_i \quad \text{and} \quad \theta'_{1,2} = \theta_{1,2} + \sum_{i=1}^{g-1} f_i \sigma_i.$$

Thus, (3) is a consequence of (1) and (2). □

## 5. THE PROOF OF THE MAIN RESULT

Throughout this section, we fix an integer $g \geq 3$. The purpose of this section is to prove the following theorem.

**Theorem 5.1** ($\operatorname{char}(k) = 0$). *A $\mathbb{Q}$-divisor $a\mu + b_{irr}\delta_{irr} + \sum_{i=1}^{[g/2]} b_i\delta_i$ on $\bar{M}_g$ is nef over $\bar{M}_g^{[1]}$ if and only if the following system of inequalities hold:*

$$\begin{aligned} &a \geq \max\left\{ \frac{b_i}{4i(g-i)} \mid i = 1, \ldots, [g/2] \right\} \\ &B_0 \geq B_1 \geq B_2 \geq \cdots \geq B_{[g/2]}, \\ &B^*_{[g/2]} \geq \cdots \geq B^*_2 \geq B^*_1 \geq B^*_0, \end{aligned}$$

*where $B_0$, $B^*_0$, $B_i$ and $B^*_i$ ($i = 1, \ldots, [g/2]$) are given by*

$$B_0 = 4b_{irr}, \quad B^*_0 = \frac{4b_{irr}}{g(2g-1)}, \quad B_i = \frac{b_i}{i(2i+1)} \quad \text{and} \quad B^*_i = \frac{b_i}{(g-i)(2(g-i)+1)}.$$

*Proof.* In the following proof, we denote $\delta_i$ by $\delta_{\{i,g-i\}}$. Moreover, we set

$$\overline{\upsilon}_g = \{\{i,j\} \mid 1 \leq i,j \leq g,\ i+j = g\}.$$

For a $\mathbb{Q}$-divisor $D = a\mu + b_{irr}\delta_{irr} + \sum_{\{i,j\}\in\overline{\upsilon}_g} b_{\{i,j\}}\delta_{\{i,j\}}$, let us consider the following inequalities:

$$a \geq \frac{b_{\{s,t\}}}{4st} \quad (\forall\{s,t\} \in \overline{\upsilon}_g) \tag{5.1.1}$$

$$4b_{irr} \geq \frac{b_{\{s,t\}}}{s(2s+1)}, \quad \frac{b_{\{s,t\}}}{t(2t+1)} \geq \frac{4b_{irr}}{g(2g-1)} \quad (\forall\{s,t\} \in \overline{\upsilon}_g \text{ with } s \leq t) \tag{5.1.2}$$

$$\frac{b_{\{l,k\}}}{l(2l+1)} \geq \frac{b_{\{s,t\}}}{s(2s+1)}, \quad \frac{b_{\{s,t\}}}{t(2t+1)} \geq \frac{b_{\{l,k\}}}{k(2k+1)} \quad (\forall\{s,t\},\{l,k\} \in \overline{\upsilon}_g \text{ with } l < s \leq t < k) \tag{5.1.3}$$

$$a \geq 0, \quad b_{irr} \geq 0, \quad b_{\{s,t\}} \geq 0 \quad (\forall\{s,t\} \in \overline{\upsilon}_g) \tag{5.1.4}$$

Let $\beta : \bar{M}_{g-1,2} \to \bar{M}_g$ and $\alpha_{s,t} : \bar{M}_{s,1} \times \bar{M}_{t,1} \to \bar{M}_g$ ($\{s,t\} \in \overline{\upsilon}_g$) be the clutching maps. First, we claim the following.

**Claim 5.1.5.** *The following are equivalent.*

(1) *$\beta^*(D)$ is nef over $M_{g-1,2}$ and $\alpha^*_{s,t}(D)$ is nef over $M_{s,1} \times M_{t,1}$ for all $\{s,t\} \in \overline{\upsilon}_g$*

(2) (5.1.1), (5.1.2), (5.1.3) *and* (5.1.4) *hold.*

On $\bar{M}_{g-1,2}$, we define $\sigma$ and $\delta_i$ ($i = 1, \dots, g-1$) as in Corollary 4.3. Moreover, we set

$$\mu' = (8g-4)\lambda - (g-1)\sigma - \sum_{i=1}^{g-1} 4i(g-1-i)\delta_i,$$
$$\theta' = (g-2)g(\psi_1+\psi_2) - 12\lambda + \sigma - \sum_{i=1}^{g-1} 4i(i-1)\delta_i.$$

Then, by using (2) of Corollary 1.5.2, we can see

$$\beta^*(D) = \frac{(g-1)(g-2)(2g-1)a - 3b_{irr}}{g(g-2)(2g-1)}\mu' + \frac{ag-b_{irr}}{g(g-2)}\theta' + \frac{(g-1)(2g+1)b_{irr}}{g(2g-1)}\sigma + \sum_{i=1}^{g-1}\left(b_{\{i,g-i\}} - \frac{4i(2i+1)}{g(2g-1)}b_{irr}\right)\delta_i.$$

Thus, by Corollary 4.3, if $\beta^*(D)$ is nef over $\bar{M}_{g-1,2}$, then

$$ag \geq b_{irr} \geq 0 \tag{5.1.6}$$

$$b_{\{i,g-i\}} \geq \frac{4i(2i+1)}{g(2g-1)}b_{irr}, \quad (i = 1, \dots, g-1) \tag{5.1.7}$$

Here we set $\mu'_1 = \theta'_1 = 0$ on $\bar{M}_{1,1}$, and

$$\mu'_e = \frac{1}{e-1}\left((8e+4)\lambda - e\delta_{irr} - \sum_{l=1}^{e-1} 4l(e-l)\delta_l\right),$$
$$\theta'_e = \frac{1}{e-1}\left(4e(e-1)\psi_1 - 12\lambda + \delta_{irr} - \sum_{l=1}^{e-1} 4l(l-1)\delta_l\right)$$

on $\bar{M}_{e,1}$ ($e \geq 2$), where $\delta_l$'s are defined as in Corollary 4.2. Let us fix $\{s,t\} \in \overline{v}_g$. Then, by using (1) of Corollary 1.5.2, we can see

$$\alpha^*_{s,t}(D) = p^*(D_s) + q^*(D_t),$$

where $p : \bar{M}_{s,1} \times \bar{M}_{t,1} \to \bar{M}_{s,1}$ and $q : \bar{M}_{s,1} \times \bar{M}_{t,1} \to \bar{M}_{t,1}$ are the projections, and $D_s \in \mathrm{Pic}(\bar{M}_{s,1}) \otimes \mathbb{Q}$ and $D_t \in \mathrm{Pic}(\bar{M}_{t,1}) \otimes \mathbb{Q}$ are given by

$$D_s = \frac{4(g-1)s(2s+1)a - 3b_{\{s,t\}}}{4s(2s+1)}\mu'_s + \frac{4sta - b_{\{s,t\}}}{4s}\theta'_s + \left(b_{irr} - \frac{b_{\{s,t\}}}{4s(2s+1)}\right)\delta_{irr} + \sum_{l=1}^{s-1}\left(b_{\{l,g-l\}} - \frac{l(2l+1)}{s(2s+1)}b_{\{s,t\}}\right)\delta_l$$

and

$$D_t = \frac{4(g-1)t(2t+1)a - 3b_{\{s,t\}}}{4t(2t+1)}\mu'_t + \frac{4sta - b_{\{s,t\}}}{4t}\theta'_s + \left(b_{irr} - \frac{b_{\{s,t\}}}{4t(2t+1)}\right)\delta_{irr} + \sum_{l=1}^{t-1}\left(b_{\{l,g-l\}} - \frac{l(2l+1)}{t(2t+1)}b_{\{s,t\}}\right)\delta_l.$$

Thus, by using Corollary 4.2 and Lemma 1.1.4, if $\alpha^*_{s,t}(D)$ is nef over $M_{s,1} \times M_{t,1}$, then

(5.1.8) $$4sta \geq b_{\{s,t\}}$$

(5.1.9) $$b_{irr} \geq \frac{b_{\{s,t\}}}{4s(2s+1)}, \quad b_{irr} \geq \frac{b_{\{s,t\}}}{4t(2t+1)}$$

(5.1.10) $$b_{\{l,g-l\}} \geq \frac{l(2l+1)}{s(2s+1)}b_{\{s,t\}} \quad (l = 1, \ldots, s-1)$$

(5.1.11) $$b_{\{l,g-l\}} \geq \frac{l(2l+1)}{t(2t+1)}b_{\{s,t\}} \quad (l = 1, \ldots, t-1)$$

Therefore, (1) implies (5.1.6) – (5.1.11). Conversely, we assume (5.1.6) – (5.1.11). Then by using(5.1.6) and (5.1.7), we can see (5.1.4). Thus, we have

$$ag - b_{irr} \geq 0 \quad \Longrightarrow \quad (g-1)(g-2)(2g-1)a - 3b_{irr} \geq 0$$
$$4sta \geq b_{\{s,t\}} \quad \Longrightarrow \quad 4(g-1)s(2s+1)a \geq 3b_{\{s,t\}} \text{ and } 4(g-1)t(2t+1)a \geq 3b_{\{s,t\}}.$$

Therefore, by Corollary 4.2, Corollary 4.3 and Lemma 1.1.4, we can see that $\beta^*(D)$ is nef over $M_{g-1,2}$ and $\alpha^*_{s,t}(D)$ is nef over $M_{s,1} \times M_{t,1}$ for all $\{s,t\} \in \overline{v}_g$. Hence it is sufficient to see that the system of inequalities (5.1.6) – (5.1.11) is equivalent to (5.1.1) – (5.1.3) under the assumption (5.1.4).

The case $s = 1, t = g-1$ in (5.1.8) and the case $i = g-1$ in (5.1.7) produce inequalities

$$4(g-1)a \geq b_{\{1,g-1\}} \quad \text{and} \quad b_{\{1,g-1\}} \geq \frac{4(g-1)}{g}b_{irr}$$

respectively, which gives rise to (5.1.6). Moreover, it is easy to see that (5.1.7) and (5.1.9) are equivalent to (5.1.2), so that it is sufficient to see that (5.1.10) and (5.1.11) are equivalent to (5.1.3).

From now on , we assume $s \leq t$. Since $s(2s+1) \leq t(2t+1)$, (5.1.10) and (5.1.11) are equivalent to saying that

(5.1.12) $$\frac{b_{\{l,k\}}}{l(2l+1)} \geq \frac{b_{\{s,t\}}}{s(2s+1)} \quad (1 \leq l < s)$$

(5.1.13) $$\frac{b_{\{l,k\}}}{l(2l+1)} \geq \frac{b_{\{s,t\}}}{t(2t+1)} \quad (s < l < t),$$

where $k = g - l$. In (5.1.12), $t < k \leq g-1$, Thus, (5.1.12) is nothing more than

$$\frac{b_{\{l,k\}}}{l(2l+1)} \geq \frac{b_{\{s,t\}}}{s(2s+1)} \quad (1 \leq l < s \leq t < k \leq g-1)$$

Moreover, in (5.1.13), $s < k < t$. Thus, (5.1.13) is nothing more that

$$\frac{b_{\{l,k\}}}{k(2k+1)} \geq \frac{b_{\{s,t\}}}{t(2t+1)} \quad (1 \leq s < l \leq k < t \leq g-1).$$

Therefore, replacing $\{s,t\}$ and $\{l,k\}$, we have

$$\frac{b_{\{s,t\}}}{t(2t+1)} \geq \frac{b_{\{l,k\}}}{k(2k+1)} \quad (1 \leq l < s \leq t < k \leq g-1).$$

Thus, we get Claim 5.1.5.

By Claim 5.1.5, it is sufficient to show the following claim to complete the proof of Theorem 5.1.

**Claim 5.1.14.** (1) *$D$ is nef over $\bar{M}_g^{[1]}$ if and only if $D$ is nef over $M_g$, $\beta^*(D)$ is nef over $M_{g-1,2}$, and $\alpha_{s,t}^*(D)$ is nef over $M_{s,1} \times M_{t,1}$ for all $\{s,t\} \in \overline{\upsilon}_g$.*

(2) *$D$ is nef over $M_g$ if and only if* (5.1.4) *holds.*

(3) (5.1.1), (5.1.2) *and* (5.1.3) *imply* (5.1.4)

(1) is obvious because

$$\bar{M}_g^{[1]} = M_g \cup \beta_g(M_{g-1,2}) \cup \bigcup_{\{s,t\} \in \overline{\upsilon}_g} \alpha_{s,t}(M_{s,1} \times M_{t,1}).$$

(2) is a consequence of [11, Theorem C]. For (3), let us consider the case $s = 1$, $t = g-1$ in (5.1.2). Then, we have

$$12 b_{irr} \geq b_{\{1,g-1\}} \quad \text{and} \quad b_{\{1,g-1\}} \geq \frac{4(g-1)}{g} b_{irr},$$

which imply $b_{irr} \geq 0$. Thus, we can see (5.1.4) using (5.1.1) and (5.1.2). □

**Corollary 5.2** ($\mathrm{char}(k) = 0$)**.** *Let $\widetilde{\Delta}_{irr}$ and $\widetilde{\Delta}_i$ ($i = 1, \ldots, [g/2]$) be the normalizations of the boundary components $\Delta_{irr}$ and $\Delta_i$ on $\bar{M}_g$, and $\rho_{irr} : \widetilde{\Delta}_{irr} \to \bar{M}_g$ and $\rho_i : \widetilde{\Delta}_i \to \bar{M}_g$ the induced morphisms. Then, a $\mathbb{Q}$-divisor $D$ on $\bar{M}_g$ is nef over $\bar{M}_g^{[1]}$ if and only if the following are satisfied:*

(1) *$D$ is weakly positive at any points of $M_g$.*

(2) *$\rho_{irr}^*(D)$ is weakly positive at any points of $\rho_{irr}^{-1}(\bar{M}_g^{[1]})$.*

(3) *$\rho_i^*(D)$ is weakly positive at any points of $\rho_i^{-1}(\bar{M}_g^{[1]})$ for all $i$.*

*Proof.* Let $\beta : \bar{M}_{g-1,2} \to \bar{M}_g$ be the clutching map. Then, there is a finite and surjective morphism $\beta' : \bar{M}_{g-1,2} \to \widetilde{\Delta}_{irr}$ with $\beta = \rho_{irr} \cdot \beta'$. Further, for $1 \leq i \leq [g/2]$, let $\alpha_{i,g-i} : \bar{M}_{i,1} \times \bar{M}_{g-i,1} \to \bar{M}_g$ be the clutching map. Then, there is a finite and surjective morphism $\alpha'_{i,g-i} : \bar{M}_{i,1} \times \bar{M}_{g-i,1} \to \widetilde{\Delta}_i$ with $\alpha_{i,g-i} = \rho_i \cdot \alpha'_{i,g-i}$. In particular, $\widetilde{\Delta}_{irr}$ and $\widetilde{\Delta}_i$'s are $\mathbb{Q}$-factorial. Therefore, if $D$ satisfies (1), (2) and (3), then $D$ is nef over $\bar{M}_g^{[1]}$.

Conversely, we assume that $D$ is nef over $\bar{M}_g^{[1]}$. (1) is nothing more than [11, Theorem C]. As in Theorem 5.1, we set $D = a\mu + b_{irr}\delta_{irr} + \sum_{i=1}^{[g/2]} b_i \delta_i$ on $\bar{M}_g$. If we trace-back the proof of Theorem 5.1, we can see that

$$\beta^*(D) \in \mathbb{Q}_+\mu' + \mathbb{Q}_+\theta' + \mathbb{Q}_+\sigma + \sum_i \mathbb{Q}_+\delta_i.$$

Here $\mu'$ and $\theta'$ are weakly positive at any points of $M_{g-1,2}$ by (1) of Corollary 4.3. Thus, so is $\beta^*(D) = \beta'^*(\rho^*_{irr}(D))$. Therefore, by virtue of (2) of Proposition 1.1.2, $\beta'_*(\beta^*(D)) = \deg(\beta')\rho^*_{irr}(D)$ is weakly positive at any points of $\rho^{-1}_{irr}(\bar{M}^{[1]}_g)$. Finally, let us consider (3). As in the proof of Theorem 5.1, there are $D_i \in \operatorname{Pic}(\bar{M}_{i,1}) \otimes \mathbb{Q}$ and $D_{g-i} \in \operatorname{Pic}(\bar{M}_{g-i,1}) \otimes \mathbb{Q}$ with $\alpha^*_{i,g-i}(D) = p^*(D_i) + q^*(D_{g-i})$, where $p : \bar{M}_{i,1} \times \bar{M}_{g-i,1} \to \bar{M}_{i,1}$ and $q : \bar{M}_{i,1} \times \bar{M}_{g-i,1} \to \bar{M}_{g-i,1}$ are the projections to the first factor and the second factor respectively. In the same way as for $\beta^*(D)$, we can see that $D_i$ (resp. $D_{g-i}$) is weakly positive at any points of $M_{i,1}$ (resp. $M_{g-i,1}$) by virtue of (1) of Corollary 4.2. Thus, by using (2) of Proposition 1.1.3, $\alpha^*_{i,g-i}(D)$ is weakly positive at any points of $M_{i,1} \times M_{g-i,1}$. Therefore, we get (3) by (2) of Proposition 1.1.2. □

**Corollary 5.3** ($\operatorname{char}(k) = 0$)**.** *With notation as in Corollary* 5.2*, if* $\rho^*_{irr}(D)$ *is nef over* $\rho^{-1}_{irr}(\bar{M}^{[1]}_g)$ *and* $\rho^*_i(D)$ *is nef over* $\rho^{-1}_i(\bar{M}^{[1]}_g)$ *for all* $i$*, then* $D$ *is nef over* $\bar{M}^{[1]}_g$*. In particular, the Mori cone of* $\bar{M}_g$ *is the convex hull spanned by curves lying on the boundary* $\bar{M}_g \setminus M_g$*, which gives rise to a special case of* [5, Proposition 3.1].

*Proof.* Let $\beta' : \bar{M}_{g-1,2} \to \widetilde{\Delta}_{irr}$ and $\alpha'_{i,g-i} : \bar{M}_{i,1} \times \bar{M}_{g-i,1} \to \widetilde{\Delta}_i$ be the same as in Corollary 5.2. By our assumption, $\beta^*(D) = \beta'^*(\rho^*_{irr}(D))$ is nef over $M_{g-1,2}$ and $\alpha^*_{i,g-i}(D) = {\alpha'}^*_{i,g-i}(\rho^*_i(D))$ is nef over $M_{i,1} \times M_{g-i,1}$ for every $i$. Therefore, by Claim 5.1.5 in Theorem 5.1, we can see that $D$ is nef over $\bar{M}^{[1]}_g$.

Let $\operatorname{Nef}_\Delta(\bar{M}_g)$ be the dual cone of the convex hull spanned by curves on the boundary $\Delta = \bar{M}_g \setminus M_g$. In order to see the last assertion of this corollary, it is sufficient to check $\operatorname{Nef}_\Delta(\bar{M}_g) = \operatorname{Nef}(\bar{M}_g)$, which is a consequence of the first assertion. □

**Example 5.4.** For example, the area of $(b_0, b_1)$ (resp. $(b_0, b_1, b_2)$) with $\lambda - b_0\delta_0 - b_1\delta_1$ (resp. $\lambda - b_0\delta_0 - b_1\delta_1 - b_2\delta_2$) nef over $\bar{M}^{[1]}_3$ (resp. $\bar{M}^{[1]}_4$) is the inside of the following triangle (resp.

polyhedron):



The area of $(b_0, b_1)$ with
$\lambda - b_0\delta_0 - b_1\delta_1$ nef over $\bar{M}_3^{[1]}$


$(\frac{1}{9}, \frac{1}{3}, \frac{4}{9})$
$(0,0,\frac{2}{7})$
$(\frac{1}{12},0,\frac{2}{7})$
$(\frac{1}{15}, \frac{1}{5},0)$
$(\frac{1}{10}, \frac{1}{5},0)$
$(0,0,0)$
$(\frac{1}{12},0,0)$

The area of $(b_0, b_1, b_2)$ with
$\lambda - b_0\delta_0 - b_1\delta_1 - b_2\delta_2$ nef over $\bar{M}_4^{[1]}$

## REFERENCES

[1] M. Cornalba and J. Harris, Divisor classes associated to families of stable varieties, with application to the moduli space of curves, Ann. Scient. Ec. Norm. Sup., 21 (1988), 455–475.
[2] P. Deligne and D. Mumford, The irreducibility of the space of curves of given genus, Publ. Math. IHES, 36 (1969), 75–110.
[3] A. J. De Jong, Smoothness, semi-stability and alteration, Publ. Math. IHES, 83 (1996), 51–93.
[4] C. Faber, Intersection-theoretical computation on $\bar{M}_g$, Banach Center Publ. 36, Polish Acad. Sci., Warsaw, 71–81. (alg-geom/9504005).
[5] A. Gibney, S. Keel and I. Morrison, Towards the ample cone of $\bar{M}_{g,n}$, (math.AG/0006208).
[6] R. Hain, Moriwaki's inequality and generalizations, talk at Chicago, http://www.math.duke.edu/~hain/talks/
[7] S. Keel and J. McKernan, Contractible extremal rays on $\bar{M}_{0,n}$, preprint, (alg-geom/9607009).
[8] J. Kollár and S. Mori, Birational geometry of algebraic varieties, Cambridge tracts in Mathematics, 134.
[9] F. Knudsen, The projectivity of the moduli space of stable curves, II and III, Math. Scand. 52 (1983), 161–199 and 200–212.
[10] A. Moriwaki, A sharp slope inequality for general stable fibrations of curves, J. reine angew. Math. 480 (1996), 177–195.
[11] A. Moriwaki, Relative Bogomolov's inequality and the cone of positive divisors on the moduli space of stable curves, J. of AMS, 11 (1998), 569–600.
[12] A. Moriwaki, The continuity of Deligne's pairing, International Mathematics Research Notices, (1999), No.19, 1057–1066.
[13] C. S. Seshadri, Quotient spaces modulo reductive algebraic groups, Ann. of Math., 95 (1972), 511–556.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KYOTO UNIVERSITY, KYOTO, 606-8502, JAPAN
*E-mail address*: `moriwaki@kusm.kyoto-u.ac.jp`